arXiv:q-alg/9709026v2 29 Sep 1997

THE QUANTIZED KNIZHNIK-ZAMOLODCHIKOV EQUATION IN TENSOR PRODUCTS OF IRREDUCIBLE sl_2-MODULES

E. MUKHIN AND A. VARCHENKO

**Department of Mathematics, University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA*
E-mail addresses: mukhin@math.unc.edu, av@math.unc.edu

August, 1997

Abstract. We consider the quantized Knizhnik-Zamolodchikov difference equation (qKZ) with values in a tensor product of irreducible sl_2 modules, the equation defined in terms of rational R-matrices. We solve the equation in terms of multidimensional q-hypergeometric integrals. We identify the space of solutions of the qKZ equation with the tensor product of the corresponding modules over the quantum group $U_q sl_2$. We compute the monodromy of the qKZ equation in terms of the trigonometric R-matrices.

1. Introduction.

1.1. In this paper we solve the rational quantized Knizhnik-Zamolodchikov difference equation (qKZ) with values in a tensor product of irreducible highest weight sl_2-modules. The rational qKZ equation is a system of difference equations for a function $\Psi(z_1, ..., z_n)$ with values in a tensor product $M_1 \otimes ... \otimes M_n$ of sl_2 modules. The system of equations has the form

$$\Psi(z_1, \dots, z_m + p, \dots, z_n) = R_{m,m-1}(z_m - z_{m-1} + p) \dots R_{m,1}(z_m - z_1 + p) e^{-\mu h_m} \times$$
$$\times R_{m,n}(z_m - z_n) \dots R_{m,m+1}(z_m - z_{m+1}) \Psi(z_1, \dots, z_n),$$

$m = 1, \dots, n$, where p, μ are complex parameters of the qKZ equation, h is a generator of the Cartan subalgebra of sl_2, h_m is the operator h acting in the m-th factor, $R_{i,j}(x)$ is the rational R-matrix $R_{M_i M_j}(x) \in \mathrm{End}(M_i \otimes M_j)$ acting in the i-th and j-th factors of the tensor product. In this paper we consider only steps p with negative real part.

The qKZ equation is an important system of difference equations. The qKZ equation was introduced in [FR] as an equation for matrix elements of vertex operators of a quantum affine algebra. An important special case of the qKZ equation had been introduced earlier in [S] as equations for form factors in integrable quantum field theory. Later, the qKZ equation was derived as an equation for correlation functions in lattice integrable models, cf. [JM] and references therein.

Solutions of the rational qKZ equation with values in a tensor product of sl_2 Verma modules $V_{\lambda_1} \otimes ... \otimes V_{\lambda_n}$ with generic highest weights $\lambda_1, \dots, \lambda_n$ were constructed in [TV1]. The solutions have the form

$$\Psi(z) = \sum_{k_1, \dots, k_n} I_{k_1, \dots, k_n}(z) f^{k_1} v_1 \otimes ... \otimes f^{k_n} v_n,$$

where $\{f^{k_1}v_1\otimes\ldots\otimes f^{k_n}v_n\}\in V_{\lambda_1}\otimes...\otimes V_{\lambda_n}$ is the standard basis in the tensor product of sl_2 Verma modules, and the coefficients $I_{k_1,\ldots,k_n}(z)$ are given by suitable multidimensional q-hypergeometric integrals.

The space of solutions of the qKZ equation with values in a tensor product of sl_2 Verma modules with generic highest weights was described in [TV1] in terms of the representation theory of the quantum group $U_q(sl_2)$ with $q=e^{\pi i/p}$. Namely, consider the tensor product $V^q_{\lambda_1}\otimes...\otimes V^q_{\lambda_n}$ of $U_q(sl_2)$ Verma modules, where $V^q_{\lambda_j}$ is the deformation of the sl_2 Verma module V_{λ_j}. It was shown in [TV1] that there is a natural isomorphism of the space $\mathcal{S}$ of meromorphic solutions of the qKZ equation with values in the tensor product $V_{\lambda_1}\otimes...\otimes V_{\lambda_n}$ and the space $V^q_{\lambda_1}\otimes...\otimes V^q_{\lambda_n}\otimes F$, where F is the space of meromorphic functions in $z_1,...,z_n$, p-periodic with respect to each of the variables,

$$V^q_{\lambda_1}\otimes...\otimes V^q_{\lambda_n}\otimes F\ \simeq\ \mathcal{S}.$$

This isomorphism was used in [TV1] to compute asymptotic solutions of the qKZ equation with values in a tensor product of sl_2 Verma modules with generic highest weights and to compute the transition functions between asymptotic solutions in terms of the trigonometric R-matrices acting in $V^q_{\lambda_1}\otimes...\otimes V^q_{\lambda_n}$.

Assume that the Verma module V_{λ_j} is reducible and vectors $\{f^kv_j\}_{k\geq N_j}$ generate a proper submodule $S_{\lambda_j}\subset V_{\lambda_j}$. Then the vectors $\{f^kv_j\}_{k<N_j}$ form a basis in the irreducible module $L_{\lambda_j}=V_{\lambda_j}/S_{\lambda_j}$. If V_{λ_j} is irreducible, then we set $N_j=\infty$.

Return to the sum $\Psi(z)$. If the Verma modules of the tensor product $V_{\lambda_1}\otimes...\otimes V_{\lambda_n}$ become reducible, then some of the coefficients in the sum become divergent. In this paper we show that the restricted sum

$$\Psi^0(z)\ =\sum_{k_1<N_1,\ldots,k_n<N_n} I_{k_1,\ldots,k_n}(z)f^{k_1}v_1\otimes\ldots\otimes f^{k_n}v_n$$

remains well defined even when some of the Verma modules become reducible.

Moreover, we show that the sum $\Psi^0(z)$ defines a solution of the qKZ equation with values in the tensor product $L_{\lambda_1}\otimes...\otimes L_{\lambda_n}$ of irreducible sl_2 modules, and under certain conditions all solutions have this form.

These results allow us to describe the space of solutions to the qKZ equation with values in $L_{\lambda_1}\otimes\ldots\otimes L_{\lambda_n}$ in terms of representation theory of the quantum group $U_q(sl_2)$. Namely, consider the tensor product $L^q_{\lambda_1}\otimes...\otimes L^q_{\lambda_n}$ of U_qsl_2 modules, where $L^q_{\lambda_j}$ is the deformation of the sl_2 irreducible module L_{λ_j}. We show that there is a natural isomorphism of the space $\mathcal{S}$ of meromorphic solutions of the qKZ equation with values in the tensor product $L_{\lambda_1}\otimes...\otimes L_{\lambda_n}$ and the space $L^q_{\lambda_1}\otimes...\otimes L^q_{\lambda_n}\otimes F$,

$$L^q_{\lambda_1}\otimes...\otimes L^q_{\lambda_n}\otimes F\ \simeq\ \mathcal{S}.$$

We compute asymptotic solutions of the qKZ equation with values in a tensor product of irreducible sl_2 Verma modules and the transition functions between the asymptotic solutions. The transition functions are given in terms of the trigonometric R-matrices acting in $L^q_{\lambda_1}\otimes...\otimes L^q_{\lambda_n}$.

In this paper we consider the rational qKZ equation associated with sl_2. There are other types of the qKZ equation: the trigonometric qKZ equation [FR], [TV2] and the elliptic qKZB equation [F1],[F2], [FTV1], [FTV2]. The trigonometric qKZ equation with values in a tensor product of $U_q(sl_2)$ Verma modules with generic highest weights and the elliptic qKZB equation with values in a tensor product of $E_{\tau,\eta}(sl_2)$ Verma modules with generic highest weights were solved in [TV2] and [FTV2], respectively. Here $E_{\tau,\eta}(sl_2)$ is the elliptic quantum group associated to sl_2. In the next paper we will extend our results to the trigonometric qKZ and elliptic qKZB equations and describe solutions of these equations with values in irreducible finite dimensional modules.

The paper is organized as follows.

In Section 2 we recall some facts about sl_2, $U_q(sl_2)$ and their representations. We define the rational R-matrix and the qKZ equation. In Section 3 we describe integral representations of solutions of the qKZ equation with values in a tensor product of sl_2 Verma modules. The statements of results are given in Section 4. The proofs are collected in Section 5.

The authors thank V.Tarasov for useful discussions.

2. General definitions and notations

2.1. **The Lie algebra sl_2.** Let e, f, h be generators of the Lie algebra sl_2 such that

$$[h,e]=e, \qquad [h,f]=-f, \qquad [e,f]=2h.$$

For an sl_2 module M, let M^* be its restricted dual with an sl_2 module structure defined by

$$\langle e\varphi, x\rangle = \langle \varphi, fx\rangle, \qquad \langle f\varphi, x\rangle = \langle \varphi, ex\rangle, \qquad \langle h\varphi, x\rangle = \langle \varphi, hx\rangle$$

for all $x\in M$, $\varphi\in M^*$. The module M^* is called the *dual* module.

For $\lambda\in\mathbb{C}$, denote V_λ the sl_2 Verma module with highest weight λ. Then $V_\lambda = \bigoplus_{i=0}^{\infty}\mathbb{C}f^iv$, where v is a highest weight vector. Denote L_λ the irreducible module with highest weight λ.

Let $\Lambda^+ = \{0,\frac12,1,\frac32,2,...\}$ be the set of dominant weights. If $\lambda\in\Lambda^+$, then L_λ is a $(2\lambda+1)$-dimensional module and

$$L_\lambda \simeq V_\lambda/S_\lambda,$$

where $S_\lambda = \bigoplus_{i=2\lambda+1}^{\infty}\mathbb{C}f^iv \subset V_\lambda$ is the maximal proper submodule. The vectors f^iv, $i=0,\dots,2\lambda$, generate a basis in L_λ.

For $\lambda\notin\Lambda^+$, $L_\lambda = V_\lambda$. It is convenient to introduce S_λ to be the zero submodule of V_λ, then $L_\lambda\simeq V_\lambda/S_\lambda$, as we have for $\lambda\in\Lambda^+$.

For an sl_2 module M with highest weght λ, denote by $(M)_l$ the subspace of weight $\lambda-l$, by $(M)^{sing}$ the kernel of the operator e, and by $(M)_l^{sing}$ the subspace $(M)_l\bigcap(M)^{sing}$.

2.2. **The algebra U_qsl_2.** Let q be a complex number different from ±1. Let e_q, f_q, q^h, q^{-h} be generators of U_qsl_2 such that

$$q^hq^{-h}=q^{-h}q^h=1, \qquad [e_q,f_q]=\frac{q^{2h}-q^{-2h}}{q-q^{-1}},$$

$$q^he_q=qe_qq^h, \qquad q^hf_q=q^{-1}f_qq^h.$$

A comultiplication $\Delta : U_qsl_2\to U_qsl_2\otimes U_qsl_2$ is given by

$$\Delta(q^h)=q^h\otimes q^h, \qquad \Delta(q^{-h})=q^{-h}\otimes q^{-h},$$

$$\Delta(e_q)=e_q\otimes q^h+q^{-h}\otimes e_q, \qquad \Delta(f_q)=f_q\otimes q^h+q^{-h}\otimes f_q.$$

The comultiplication defines a module structure on tensor products of U_qsl_2 modules.

For $\lambda \in \mathbb{C}$, denote V^q_λ the $U_q sl_2$ Verma module with highest weight q^λ. Then $V^q_\lambda = \bigoplus_{i=0}^{\infty} \mathbb{C} f_q^i v^q$, where v^q is a highest weight vector.

For $\lambda \in \Lambda^+$, $S^q_\lambda = \bigoplus_{i=2\lambda+1}^{\infty} \mathbb{C} f_q^i v^q$ is a submodule in V^q_λ. Denote L^q_λ the quotient module $V^q_\lambda / S^q_\lambda$. The module L^q_λ is the $(2\lambda+1)$-dimensional highest weight module with highest weight q^λ. The vectors $f_q^i v^q$, $i = 0, \dots, 2\lambda$, generate a basis in L^q_λ.

For $\lambda \notin \Lambda^+$, let $L^q_\lambda = V^q_\lambda$. It is convenient to introduce S^q_λ to be the zero submodule of V^q_λ, then $L^q_\lambda \simeq V^q_\lambda / S^q_\lambda$, as we have for $\lambda \in \Lambda^+$.

For an $U_q sl_2$ module M^q with highest weight q^λ, denote by $(M^q)_l$ the subspace of weight $q^{\lambda - l}$, by $(M^q)^{sing}$ the kernel of the operator e_q, and by $(M^q)^{sing}_l$ the subspace $(M^q)_l \bigcap (M^q)^{sing}$.

2.3. **The rational R-matrix.** The *Yangian* $Y(sl_2)$ is a Hopf algebra which has a family of homomorphisms to the universal enveloping algebra of sl_2, $Y(sl_2) \to U(sl_2)$, depending on a complex parameter. Therefore, each sl_2 module M carries a Yangian module structure $M(x)$ depending on a parameter, see [CP],[TV1].

For sl_2 irreducible highest weight modules $L_{\lambda_1}, L_{\lambda_2}$ and generic numbers $x, y \in \mathbb{C}$, the Yangian modules $L_{\lambda_1}(x) \otimes L_{\lambda_2}(y)$ and $L_{\lambda_2}(y) \otimes L_{\lambda_1}(x)$ are irreducible and isomorphic. There exists a unique intertwiner which sends $v_1 \otimes v_2$ to $v_2 \otimes v_1$, where v_i is a highest vector in L_{λ_i}, $i = 1, 2$. This intertwiner has the form $PR_{L_{\lambda_1} L_{\lambda_2}}(x - y)$, where P is the operator of permutation of factors. The operator $R_{L_{\lambda_1} L_{\lambda_2}}(x) \in \mathrm{End}(L_{\lambda_1} \otimes L_{\lambda_2})$ is called the *rational R-matrix*, see details in [TV1]. The rational R-matrix commutes with the sl_2 action on the tensor product $L_{\lambda_1} \otimes L_{\lambda_2}$.

Let $\lambda_1, \lambda_2 \notin \Lambda^+$. Let

$$V_{\lambda_1} \otimes V_{\lambda_2} = \bigoplus_{l=0}^{\infty} V_{\lambda_1 + \lambda_2 - l}$$

be the decomposition of the tensor product of sl_2 Verma modules into the direct sum of irreducibles, and let $\Pi^{(l)}$ be the projector on $V_{\lambda_1 + \lambda_2 - l}$ along the other summands.

There is a formula for the rational R-matrix,

$$R_{V_{\lambda_1} V_{\lambda_2}}(x) = \sum_{l=0}^{\infty} \Pi^{(l)} \prod_{s=0}^{l-1} \frac{x + \lambda_1 + \lambda_2 - s}{x - \lambda_1 - \lambda_2 + s},$$

see [KRS],[T].

Let $\lambda_1, \lambda_2 \in \Lambda^+$. Let

$$L_{\lambda_1} \otimes L_{\lambda_2} = \bigoplus_{l=0}^{2\min\{\lambda_1, \lambda_2\}} L_{\lambda_1 + \lambda_2 - l}$$

be the decomposition of the tensor product of finite dimensional irreducible sl_2 modules into the direct sum of irreducibles, and let $\Pi^{(l)}$ be the projector on $L_{\lambda_1 + \lambda_2 - l}$ along the other summands.

There is a formula for the rational R-matrix,

$$R_{L_{\lambda_1} L_{\lambda_2}}(x) = \sum_{l=0}^{2\min\{\lambda_1, \lambda_2\}} \Pi^{(l)} \prod_{s=0}^{l-1} \frac{x + \lambda_1 + \lambda_2 - s}{x - \lambda_1 - \lambda_2 + s},$$

see [KRS],[T].

The vector spaces $V_{\lambda_1}\otimes V_{\lambda_2}$ for different values of λ_1,λ_2 are identified by distinguished bases $\{f^{l_1}v_1\otimes f^{l_2}v_2\,|\,l_1,l_2\in\mathbb{Z}_{\geq 0}\}$.

Theorem 1. *1. The rational R-matrix* $R_{V_{\lambda_1}V_{\lambda_2}}(x)\in \mathrm{End}(V\otimes V)$ *is a meromorphic function of* x,λ_1,λ_2. *The poles of* $R_{V_{\lambda_1}V_{\lambda_2}}(x)$ *have the form* $x-\lambda_1-\lambda_2+s=0$, *where* $s\in\mathbb{Z}_{\geq 0}$.

2. Let x *be generic. Then the rational R-matrix* $R_{V_{\lambda_1}V_{\lambda_2}}(x)$ *preserves* $S_{\lambda_1}\otimes V_{\lambda_2}+V_{\lambda_1}\otimes S_{\lambda_2}$.

3. Let $V_{\lambda_1}\otimes V_{\lambda_2}\to L_{\lambda_1}\otimes L_{\lambda_2}$ *be the canonical factorization map. Then for generic* x, *the rational R-matrix* $R_{V_{\lambda_1}V_{\lambda_2}}(x)$ *can be factorized to an operator* $R(x)\in\mathrm{End}(L_{\lambda_1}\otimes L_{\lambda_2})$ *and, moreover,* $R(x)=R_{L_{\lambda_1}L_{\lambda_2}}(x)$.

An analog of Theorem 1 for the elliptic R-matrix is proved in Theorems 8, 31 in [FTV1]. The same proof works for the rational R-matrix. We give an alternative proof in Section 5.

2.4. **The trigonometric R-matrix.** Let q be a complex number, and not a root of unity. The *quantum affine algebra* $\widehat{U_q sl_2}$ is a Hopf algebra which has a family of homomorphisms, $\widehat{U_q sl_2}\to U_q sl_2$, depending on a complex parameter. Therefore, each $U_q sl_2$ module M^q carries a $\widehat{U_q sl_2}$ module structure $M^q(x)$ depending on a parameter, see [T], [CP].

For $U_q sl_2$ irreducible highest weight modules $L^q_{\lambda_1}, L^q_{\lambda_2}$ and generic numbers $x,y\in\mathbb{C}$, the $\widehat{U_q sl_2}$ modules $L^q_{\lambda_1}(x)\otimes L^q_{\lambda_2}(y)$ and $L^q_{\lambda_2}(y)\otimes L^q_{\lambda_1}(x)$ are isomorphic. There exists a unique intertwiner which sends $v^q_1\otimes v^q_2$ to $v^q_2\otimes v^q_1$, where v^q_i is a highest vector in $L^q_{\lambda_i}$, $i=1,2$. This intertwiner has the form $PR^q_{L^q_{\lambda_1}L^q_{\lambda_2}}(x/y)$, where P is the operator of permutation of factors. The operator $R^q_{L^q_{\lambda_1}L^q_{\lambda_2}}(x)\in\mathrm{End}(L^q_{\lambda_1}\otimes L^q_{\lambda_2})$ is called the *trigonometric R-matrix*, see details in [TV1]. The trigonometric R-matrix preserves the weight decomposition.

Let $\lambda_1,\lambda_2\notin\Lambda^+$. Let

$$V^q_{\lambda_1}\otimes V^q_{\lambda_2}=\bigoplus_{l=0}^{\infty}V^q_{\lambda_1+\lambda_2-l}$$

be the decomposition of the tensor product of $U_q sl_2$ Verma modules into the direct sum of irreducible modules, and let $\Pi^{(l)}$ be the projector on $V^q_{\lambda_1+\lambda_2-l}$ along the other summands.

There is a formula for the trigonometric R-matrix:

$$R^q_{V^q_{\lambda_1}V^q_{\lambda_2}}(x)=R^q_{\lambda_1,\lambda_2}(0)\sum_{l=0}^{\infty}\Pi^{(l)}\prod_{s=0}^{l-1}\frac{1-xq^{2s-2\lambda_1-2\lambda_2}}{1-xq^{2\lambda_1+2\lambda_2-2s}},$$

where

$$R^q_{\lambda_1,\lambda_2}(0)=q^{2\lambda_1\lambda_2-2h\otimes h}\sum_{k=0}^{\infty}(q^2-1)^{2k}\prod_{s=1}^{k}(1-q^{2s})^{-1}\,(q^h f_q\otimes q^{-h}e_q)^k,$$

see [CP],[T].

Let $\lambda_1,\lambda_2\in\Lambda^+$. Let

$$L^q_{\lambda_1}\otimes L^q_{\lambda_2}=\bigoplus_{l=0}^{2\min\{\lambda_1,\lambda_2\}}L^q_{\lambda_1+\lambda_2-l}$$

be the decomposition of the tensor product of finite dimensional irreducible $U_q sl_2$ modules into the direct sum of irreducibles, and let $\Pi^{(l)}$ be the projector on $L^q_{\lambda_1+\lambda_2-l}$ along the other summands.

There is a formula for the trigonometric R-matrix,

$$R^q_{L^q_{\lambda_1} L^q_{\lambda_2}}(x) = R^q_{\lambda_1,\lambda_2}(0) \sum_{l=0}^{2\min\{\lambda_1,\lambda_2\}} \Pi^{(l)} \prod_{s=0}^{l-1} \frac{1 - xq^{2s-2\lambda_1-2\lambda_2}}{1 - xq^{2\lambda_1+2\lambda_2-2s}},$$

see [CP],[T]. The vector spaces $V^q_{\lambda_1} \otimes V^q_{\lambda_2}$ for different values of λ_1, λ_2 are identified by distinguished bases $\{f^{l_1} v^q_1 \otimes f^{l_2} v^q_2 \,|\, l_1, l_2 \in \mathbb{Z}_{\geq 0}\}$.

Theorem 2. *Let $q \in \mathbb{C}$ be not a root of unity.*

1. The trigonometric R-matrix $R^q_{V^q_{\lambda_1} V^q_{\lambda_2}}(x) \in \mathrm{End}(V \otimes V)$ is a meromorphic function of x, λ_1, λ_2. The poles of $R^q_{V^q_{\lambda_1} V^q_{\lambda_2}}(x)$ have the form $x = q^{-2\lambda_1-2\lambda_2+2s}$, where $s \in \mathbb{Z}_{\geq 0}$.

2. Let x be generic. Then the trigonometric R-matrix $R^q_{V^q_{\lambda_1} V^q_{\lambda_2}}(x)$ preserves $S^q_{\lambda_1} \otimes V^q_{\lambda_2} + V^q_{\lambda_1} \otimes S^q_{\lambda_2}$.

3. Let $V^q_{\lambda_1} \otimes V^q_{\lambda_2} \to L^q_{\lambda_1} \otimes L^q_{\lambda_2}$ be the canonical factorization map. Then for generic x, the trigonometric R-matrix $R^q_{V^q_{\lambda_1} V^q_{\lambda_2}}(x)$ can be factorized to an operator $R^q(x) \in \mathrm{End}(L^q_{\lambda_1} \otimes L^q_{\lambda_2})$ and, moreover, $R^q(x) = R^q_{L^q_{\lambda_1} L^q_{\lambda_2}}(x)$.

An analog of Theorem 1 for the elliptic R-matrix is proved in Theorems 8, 31 in [FTV1]. The same proof works for the trigonometric R-matrix.

2.5. **The qKZ equation.** The *rational quantized Knizhnik-Zamolodchikov equation(qKZ)* associated to sl_2 is the following system of linear difference equations for a function $\Psi(z_1, \dots, z_n)$ with values in a tensor product $M_1 \otimes \dots \otimes M_n$ of sl_2 modules:

$$\begin{aligned}\Psi(z_1, \dots, z_m + p, \dots, z_n) = R_{M_m,M_{m-1}}(z_m - z_{m-1} + p) \dots R_{M_m,M_1}(z_m - z_1 + p) e^{-\mu h_m} \times \\ \times R_{M_m,M_n}(z_m - z_n) \dots R_{M_m,M_{m+1}}(z_m - z_{m+1}) \Psi(z_1, \dots, z_n),\end{aligned}$$

for $m = 1, \dots, n$. Here p, μ are complex parameters, μ is chosen so that $0 \leq \mathrm{Im}\,\mu < 2\pi$; h_m is the operator $h \in sl_2$ acting in the m-th representation, $R_{M_i M_j}(x) \in \mathrm{End}(M_i \otimes M_j)$ is the rational R-matrix acting in the i-th and j-th factors, see [FR]. The linear operators in the right hand side of the equations are called the *qKZ operators.*

The qKZ operators commute with the action of the operator $h \in sl_2$ in the tensor product $M_1 \otimes \dots \otimes M_n$. Therefore, in order to construct all solutions of the qKZ equation, it is enough to solve the qKZ equation with values in weight spaces $(M_1 \otimes \dots \otimes M_n)_l$.

If the parameter μ of the equation is equal to zero, then the qKZ operators commute with the sl_2 action in the tensor product $M_1 \otimes \dots \otimes M_n$, and in order to construct all solutions of the qKZ equation in this case, it is enough to solve the equation with values in singular weight spaces $(M_1 \otimes \dots \otimes M_n)^{sing}_l$.

Let $\pi : V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n} \to L_{\lambda_1} \otimes \dots \otimes L_{\lambda_n}$ be the canonical projection map.

Lemma 3. *Let $\Psi(z)$ be a solution of the qKZ equation with values in $V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n}$. Then $\pi \circ \Psi(z)$ is a solution of the qKZ equation with values in $L_{\lambda_1} \otimes \dots \otimes L_{\lambda_n}$.*

Lemma 3 follows from Theorem 1.

3. The Hypergeometric pairing, [TV1]

3.1. **The phase function.** Let $z=(z_1,\dots,z_n)\in\mathbb{C}^n$, $\lambda=(\lambda_1,\dots,\lambda_n)\in\mathbb{C}^n$, $t=(t_1,\dots,t_l)\in\mathbb{C}^l$. The *phase function* is defined by the following formula:

$$\Phi_l(t,z,\lambda)=\exp(\mu\sum_{i=1}^{l}t_i/p)\prod_{i=1}^{n}\prod_{j=1}^{l}\frac{\Gamma((t_j-z_i+\lambda_i)/p)}{\Gamma((t_j-z_i-\lambda_i)/p)}\prod_{1\le i<j\le l}\frac{\Gamma((t_i-t_j-1)/p)}{\Gamma((t_i-t_j+1)/p)}.$$

3.2. **Actions of the symmetric group.** Let $f=f(t_1,\dots,t_l)$ be a function. For a permutation $\sigma\in\mathbb{S}^l$, define the functions $[f]^{rat}_\sigma$ and $[f]^{trig}_\sigma$ via the action of the simple transpositions $(i,i+1)\in\mathbb{S}^{\lessdot}$, $i=1,\dots,l-1$, given by

$$[f]^{rat}_{(i,i+1)}(t_1,\dots,t_l)=f(t_1,\dots,t_{i+1},t_i,\dots,t_l)\frac{t_i-t_{i+1}-1}{t_i-t_{i+1}+1},$$

$$[f]^{trig}_{(i,i+1)}(t_1,\dots,t_l)=f(t_1,\dots,t_{i+1},t_i,\dots,t_l)\frac{\sin(\pi(t_i-t_{i+1}-1)/p)}{\sin(\pi(t_i-t_{i+1}+1)/p)}.$$

If for all $\sigma\in\mathbb{S}^l$, $[f]^{rat}_\sigma=f$, we will say that the function is *symmetric with respect to the rational action.* If for all $\sigma\in\mathbb{S}^l$, $[f]^{trig}_\sigma=f$, we will say that the function is *symmetric with respect to the trigonometric action.*

This definition implies the following important Remark.

Remark. If $w(t_1,\dots,t_l)$ is $\mathbb{S}^l$ symmetric with respect to the rational action and $W(t_1,\dots,t_l)$ is $\mathbb{S}^l$ symmetric with respect to the trigonometric action, then $\Phi_l wW$ is a symmetric function of $t_1,\dots,t_l$ (in the usual sense).

3.3. **Rational weight functions.** Fix natural numbers n,l.

Set $\mathcal{Z}^n_l=\{\bar l=(l_1,\dots,l_n)\in\mathbb{Z}^n_{\ge0}\,|\,\sum\limits_{i=1}^{n}l_i=l\}$. For $\bar l\in\mathcal{Z}^n_l$ and $m=0,1,\dots,n$, set $l^m=\sum\limits_{i=1}^{m}l_i$.

For $\bar l\in\mathcal{Z}^n_l$, define the *rational weight function* $w_{\bar l}$ by

$$w_{\bar l}(t,z,\lambda)=\sum_{\sigma\in\mathbb{S}^l}\left[\prod_{m=1}^{n}\frac{1}{l_m!}\prod_{j=l^{m-1}+1}^{l^m}\left(\frac{1}{t_j-z_m-\lambda_m}\prod_{k=1}^{m}\frac{t_j-z_k+\lambda_k}{t_j-z_k-\lambda_k}\right)\right]^{rat}_\sigma.$$

For fixed $z,\lambda\in\mathbb{C}^n$, the space spanned over $\mathbb{C}$ by all rational weight functions $w_{\bar l}(t,z,\lambda)$, $\bar l\in\mathcal{Z}^n_l$, is called the *hypergeometric rational space specialized at* z,λ and is denoted $\mathfrak{F}(z,\lambda)=\mathfrak{F}^{\,n}_{\,l}(z,\lambda)$. This space is a space of functions of variable t.

For generic values of z,λ, the space $\mathfrak{F}(z,\lambda)$ can be identified with the space $(V^*_{\lambda_1}\otimes\dots\otimes V^*_{\lambda_n})_l$ by the map

$$\mathfrak{a}(z,\lambda):\ (f^{l_1}v_1)^*\otimes\dots\otimes(f^{l_n}v_n)^*\mapsto w_{\bar l}(t,z,\lambda).$$

Here $\{(f^lv_i)^*\,|\,l\in\mathbb{Z}_{\ge0}\}$ is the basis of $V^*_{\lambda_i}$, dual to the standard basis of V_{λ_i} given by $\{f^lv_i\,|\,l\in\mathbb{Z}_{\ge0}\}$, see Lemma 4.5, Corallary 4.8 in [TV1].

3.4. **Trigonometric weight functions.** Fix natural numbers n, l.

For $\bar{l} \in \mathcal{Z}_l^n$, define the *trigonometric weight function* $W_{\bar{l}}$ by

$$W_{\bar{l}}(t,z,\lambda) =$$
$$\sum_{\sigma\in\mathbb{S}^l}\left[\prod_{m=1}^{n}\prod_{s=1}^{l_m}\frac{\sin(\pi/p)}{\sin(\pi s/p)}\prod_{j=l^{m-1}+1}^{l^m}\frac{\exp(\pi i(z_m-t_j)/p)}{\sin(\pi(t_j-z_m-\lambda_m)/p)}\prod_{k=1}^{m}\frac{\sin(\pi(t_j-z_k+\lambda_k)/p)}{\sin(\pi(t_j-z_k-\lambda_k)/p)}\right]_{\sigma}^{trig}.$$

A function $W(t,z,\lambda)$ is said to be a *holomorphic trigonometric weight function* if

$$W(t,z,\lambda) = \sum_{m_1+\ldots+m_n=l} a_{\bar{m}}(\lambda, e^{2\pi i z_1/p},\ldots,e^{2\pi i z_n/p})W_{\bar{m}}(t,z,\lambda), \tag{1}$$

where $a_{\bar{m}}(\lambda,u)$ are holomorphic functions of parameters $\lambda, u \in \mathbb{C}^n$. We denote $\mathfrak{G}$ the space of all holomorphic trigonometric weight functions. This space is a space of functions of variables of t,z,λ.

For a permutation $\sigma \in \mathbb{S}^n$, define the *trigonometric weight functions* $W_{\bar{l}}^{\sigma}$ by $W_{\bar{l}}^{\sigma}(t,z,\lambda) = W_{\sigma\bar{l}}(t,\sigma z,\sigma\lambda)$, where $\sigma\bar{l} = (l_{\sigma_1},\ldots,l_{\sigma_n})$, $\sigma z = (z_{\sigma_1},\ldots,z_{\sigma_n})$, $\sigma\lambda = (\lambda_{\sigma_1},\ldots,\lambda_{\sigma_n})$.

For fixed $\lambda, z \in \mathbb{C}^n$ and $\sigma \in \mathbb{S}^n$, the space spanned over $\mathbb{C}$ by all trigonometric weight functions $W_{\bar{l}}^{\sigma}(t,z,\lambda)$, $\bar{l} \in \mathcal{Z}_l^n$, is called the *hypergeometric trigonometric space specialized at* z,λ and is denoted $\mathfrak{G}^{\,\sigma}(z,\lambda) = \mathfrak{G}_{\,l}^{\,n,\sigma}(z,\lambda)$. This space is a space of functions of variable t.

For generic z,λ, the space $\mathfrak{G}^{\,\sigma}(z,\lambda)$ does not depend on σ, see Section 2 in [TV1], we denote it $\mathfrak{G}\,(z,\lambda)$.

For $\bar{l} \in \mathcal{Z}_l^{n-1}$, define the *singular trigonometric weight function* $W_{\bar{l}}^{sing}$ by

$$W_{\bar{l}}^{sing}(t,z,\lambda) =$$
$$\sum_{\sigma\in\mathbb{S}^l}\Bigg[\prod_{m=1}^{n-1}\prod_{s=1}^{l_m}\frac{\sin(\pi/p)}{\sin(\pi s/p}\sin(\pi(z_m-\lambda_m-z_{m+1}-\lambda_{m+1}+s-1)/p)\times$$
$$\times\prod_{j=l^{m-1}+1}^{l^m}\frac{1}{\sin(\pi(t_j-z_m-\lambda_m)/p)\sin(\pi(t_j-z_{m+1}-\lambda_{m+1})/p)}\times$$
$$\times\prod_{k=1}^{m}\frac{\sin(\pi(t_j-z_k+\lambda_k)/p)}{\sin(\pi(t_j-z_k-\lambda_k)/p)}\Bigg]_{\sigma}^{trig}.$$

For fixed $z,\lambda \in \mathbb{C}^n$, the space spanned over $\mathbb{C}$ by all singular trigonometric weight functions $W_{\bar{l}}(t,z,\lambda)$, $\bar{l} \in \mathcal{Z}_l^n$, is called the *singular hypergeometric trigonometric space specialized at* z,λ and is denoted $\mathfrak{G}^{\,sing}(z,\lambda) = \mathfrak{G}_{\,l}^{\,sing,n}(z,\lambda)$.

We have $\mathfrak{G}^{\,sing}(z,\lambda) \subset \mathfrak{G}\,(z,\lambda)$, see Lemma 2.29 in [TV1].

A function $W(t,z,\lambda)$ is said to be a *holomorphic singular trigonometric weight function* if $W(t,z,\lambda)$ is a holomorphic trigonometric weight function, and for all $z,\lambda \in \mathbb{C}^n$, the function $W(t,z,\lambda)$ belongs to $\mathfrak{G}^{\,sing}(z,\lambda)$. We denote $\mathfrak{G}^{\,sing}$ the space of all holomorphic singular trigonometric weight functions. This space is a space of functions of variables t,z,λ.

Lemma 4. *For any* $\bar{l} \in \mathcal{Z}_l^{n-1}$*, the function* $W_{\bar{l}}^{sing}(t,z,\lambda)$ *belongs to* $\mathfrak{G}^{\,sing}$.

Proof: By Lemmas 2.28 and 2.29 of [TV1], we have the decomposition (1),

$$W_{\bar{l}}^{sing}(t,z,\lambda) = \sum_{m_1+\ldots+m_n=l} a_{\bar{m}}(\lambda, e^{2\pi i z_1/p}, \ldots, e^{2\pi i z_n/p}) W_{\bar{m}}(t,z,\lambda).$$

Here $a_{\bar{m}}(\lambda,u)$ are meromorphic functions of $\lambda, u \in \mathbb{C}^n$. The functions $a_{\bar{m}}(\lambda,u)$ are holomorphic for the following reason.

Suppose, for some $\bar{m}_0 \in \mathcal{Z}_l^n$ and some $Z, \Lambda \in \mathbb{C}^n$ the function $a_{\bar{m}}(\lambda,u)$ has a pole at Z, Λ. The functions $W_{\bar{m}}(t,z,\lambda)$, $\bar{m} \in \mathcal{Z}_l^n$ are linearly independent for generic t,z,λ by Lemma 2.28 in [TV1]. Moreover, for generic t, the functions $W_{\bar{m}}(t,z,\lambda)$, $\bar{m} \in \mathcal{Z}_l^n$, do not have poles at Z, Λ. Hence, for generic t, the function $W_{\bar{l}}^{sing}(t,z,\lambda)$ has a pole at Z, Λ. But this is not so. □

For $\bar{l} \in \mathcal{Z}_l^n$, define the *weight coefficient* $c_{\bar{l}}(\lambda)$ by

$$c_{\bar{l}}(\lambda) = \prod_{m=1}^{n} \prod_{s=0}^{l_m-1} \frac{\sin(\pi(s+1)/p)\sin(\pi(2\lambda_m - s)/p)}{\sin(\pi/p)}.$$

Let $q = e^{\pi i/p}$. The weight space $(V_{\lambda_{\sigma_1}}^q \otimes \ldots \otimes V_{\lambda_{\sigma_n}}^q)_l$ is mapped to the trigonometric hypergeometric space $\mathfrak{G}^{\,\sigma}(z,\lambda)$ by

$$\mathfrak{b}_\sigma(z,\lambda):\ f_q^{l_{\sigma_1}} v_{\sigma_1}^q \otimes \ldots \otimes f_q^{l_{\sigma_n}} v_{\sigma_n}^q \mapsto c_{\bar{l}}(\lambda) W_{\bar{l}}^{\sigma}(t,z,\lambda). \tag{2}$$

If z, λ are generic and q is not a root of unity, then the map $\mathfrak{b}_\sigma(z,\lambda)$ is an isomorphism of vector spaces, see Lemma 4.17 and Corallary 4.20 in [TV1]. Moreover, the singular hypergeometric space is identified with the subspace of singular vectors,

$$\mathfrak{G}^{\,sing}(z,\lambda) = \mathfrak{b}_\sigma(z,\lambda)((V_{\lambda_{\sigma_1}}^q \otimes \ldots \otimes V_{\lambda_{\sigma_n}}^q)_l^{sing}),$$

see Corallary 4.21 in [TV1].

The composition maps

$$\mathfrak{b}_{\sigma,\sigma'}(z,\lambda):\ (V_{\lambda_{\sigma'_1}}^q \otimes \ldots \otimes V_{\lambda_{\sigma'_n}}^q)_l \to (V_{\lambda_{\sigma_1}}^q \otimes \ldots \otimes V_{\lambda_{\sigma_n}}^q)_l,\ \ \mathfrak{b}_{\sigma,\sigma'}(z,\lambda) = (\mathfrak{b}_\sigma(z,\lambda))^{-1} \circ \mathfrak{b}_{\sigma'}(z,\lambda),$$

are called the *transition functions.*

Theorem 5. *(Theorem 4.22 in* [TV1]*.) Let q be not a root of unity. For any $\sigma \in \mathbb{S}^n$ and any transposition $(m, m+1), m = 1, \ldots, n-1$, the transition function*

$$\mathfrak{b}_{\sigma,\sigma\circ(m,m+1)}(z,\lambda):\ (V_{\lambda_{\sigma_1}}^q \otimes \ldots \otimes V_{\lambda_{\sigma_{m+1}}}^q \otimes V_{\lambda_{\sigma_m}}^q \otimes \ldots \otimes V_{\lambda_{\sigma_n}}^q)_l \to (V_{\lambda_{\sigma_1}}^q \otimes \ldots \otimes V_{\lambda_{\sigma_n}}^q)_l$$

equals the operator $P_{V_{\lambda_{\sigma_{m+1}}}^q V_{\lambda_{\sigma_m}}^q} R_{V_{\lambda_{\sigma_{m+1}}}^q V_{\lambda_{\sigma_m}}^q}(e^{2\pi i(z_{\sigma_{m+1}} - z_{\sigma_m})/p})$ acting in the m-th and $(m+1)$-st factors. Here $P_{V_{\lambda_{\sigma_{m+1}}}^q V_{\lambda_{\sigma_m}}^q}$ is the operator of permutation of the factors $V_{\lambda_{\sigma_{m+1}}}^q$ and $V_{\lambda_{\sigma_m}}^q$.

Corollary 6. *Let q be not a root of unity. For any $\sigma \in \mathbb{S}^n$ and any transposition $(m, m+1), m = 1, \ldots, n-1$, the transition function $\mathfrak{b}_{\sigma,\sigma'}(z,\lambda)$ can be factorized to a transition function*

$$\mathfrak{B}_{\sigma,\sigma'}(z,\lambda):\ (L_{\lambda_{\sigma'_1}}^q \otimes \ldots \otimes L_{\lambda_{\sigma'_n}}^q)_l \to (L_{\lambda_{\sigma_1}}^q \otimes \ldots \otimes L_{\lambda_{\sigma_n}}^q)_l.$$

Moreover, for any $\sigma \in \mathbb{S}^n$ *and any transposition* $(m, m+1), m = 1, \dots, n-1$, *the induced transition function*

$$\mathfrak{B}_{\sigma,\sigma\circ(m,m+1)}(z,\lambda) : \, (L^q_{\lambda_{\sigma_1}} \otimes \dots \otimes L^q_{\lambda_{\sigma_{m+1}}} \otimes L^q_{\lambda_{\sigma_m}} \otimes \dots \otimes L^q_{\lambda_{\sigma_n}})_l \to (L^q_{\lambda_{\sigma_1}} \otimes \dots \otimes L^q_{\lambda_{\sigma_n}})_l$$

equals the operator $P_{L^q_{\lambda_{\sigma_{m+1}}} L^q_{\lambda_{\sigma_m}}} R_{L^q_{\lambda_{\sigma_{m+1}}} L^q_{\lambda_{\sigma_m}}} (e^{2\pi i (z_{\sigma_{m+1}} - z_{\sigma_m})/p})$ *acting in the* m*-th and* $(m+1)$*-th factors.*

Corollary 6 follows from Theorems 2 and 5.

Thus, the map $\mathfrak{b}_\sigma(z,\lambda) : \, (V^q_{\lambda_{\sigma_1}} \otimes \dots \otimes V^q_{\lambda_{\sigma_n}})_l \to \mathfrak{G}\,(z,\lambda)$ can be factorized to a map

$$\mathfrak{B}_\sigma(z,\lambda) : \, (L^q_{\lambda_{\sigma_1}} \otimes \dots \otimes L^q_{\lambda_{\sigma_n}})_l \to \mathfrak{G}\,(z,\lambda), \tag{3}$$

defined by the same formula (2). Moreover, the image of $\mathfrak{B}_\sigma(z,\lambda)$ does not depend on σ. Transition functions $(\mathfrak{B}_\sigma(z,\lambda))^{-1} \circ \mathfrak{B}_{\sigma'}(z,\lambda)$ are well defined, and $(\mathfrak{B}_\sigma(z,\lambda))^{-1} \circ \mathfrak{B}_{\sigma'}(z,\lambda) = \mathfrak{B}_{\sigma,\sigma'}(z,\lambda)$.

3.5. **Hypergeometric integrals.**

Fix $p \in \mathbb{C}$, $\mathrm{Re}\, p < 0$. Let $\mathrm{Im}\,\mu \neq 0$. Assume that the parameters $z, \lambda \in \mathbb{C}^n$ satisfy the condition $\mathrm{Re}\,(z_i + \lambda_i) < 0$ and $\mathrm{Re}\,(z_i - \lambda_i) > 0$ for all $i = 1, \dots, n$. For a rational weight function $w_{\bar{l}}(t,z,\lambda)$, $\bar{l} \in \mathcal{Z}^n_l$, and a trigonometric weight function $W(t,z,\lambda) \in \mathfrak{G}$, define the *hypergeometric integral* $I(w,W)(z,\lambda)$ by the formula

$$I(w,W)(z,\lambda) = \int\limits_{\substack{\mathrm{Re}\, t_i = 0,\\ i=1,\dots,l}} \Phi_l(t,z,\lambda) w(t,z,\lambda) W(t,z,\lambda)\, d^l t, \tag{4}$$

where $d^l t = dt_1 \dots dt_l$.

The hypergeometric integral for generic z, λ and an arbitrary step p with negative real part is defined by analytic continuation with respect to z, λ and p. This analytic continuation makes sense since the integrand is meromorphic in z, λ and p. The poles of the integrand are located at the union of hyperplanes

$$t_i = z_k \pm (\lambda_k + sp), \qquad t_i = t_j \pm (1 - sp), \tag{5}$$

$i, j = 1, \dots, l$, $k = 1, \dots, n$, $s \in \mathbb{Z}_{\geq 0}$. We move the parameters z, λ and p in such a way that the topology of the complement in $\mathbb{C}^l$ to the union of hyperplanes (5) does not change. We deform accordingly the integration cycle (the imaginary subspace) in such a way that it does not intersect the hyperplanes (5) at any moment of the deformation. The deformed integration cycle is called the *deformed imaginary subspace* and is denoted $\mathfrak{I}\,(z,\lambda)$, $\mathfrak{I}\,(z,\lambda) \subset \mathbb{C}^l$. Then the analytic continuation of integral (4) is given by

$$I(w,W)(z,\lambda) = \int\limits_{\mathfrak{I}\,(z,\lambda)} \Phi_l(t,z,\lambda) w(t,z,\lambda) W(t,z,\lambda)\, d^l t, \tag{6}$$

see Section 5 in [TV1].

Theorem 7. *(Theorem 5.7 in* [TV1]*.) Let* $\mathrm{Im}\,\mu \neq 0$. *For any* $w_{\bar{l}}(t,z,\lambda)$, $\bar{l} \in \mathcal{Z}^n_l$, *and any* $W(t,z,\lambda) \in \mathfrak{G}$, *the hypergeometric integral* (6) *is a univalued meromorphic function of variables* p, z, λ *holomorphic on the set*

$$\mathrm{Re}\, p < 0, \qquad \{1, \dots, l\} \not\subset p\mathbb{Z},$$

$$2\lambda_m - s \notin p\mathbb{Z}, \qquad m = 1, \dots, n, \qquad s = 1 - l, \dots, l - 1, \tag{7}$$

$$z_k \pm \lambda_k - z_m \pm \lambda_m - s \notin p\mathbb{Z}, \qquad k,m = 1,\dots,n, \qquad k \neq m, \qquad s = 1-l,\dots,l-1,$$

where we allow arbitrary combinations of $\pm$*.*

The case $\mathrm{Im}\,\mu = 0$ is treated in the same manner.

Theorem 8. *(Theorem 5.8 in* [TV1]*.) Let* $\mathrm{Im}\,\mu = 0$*. For any* $w_{\bar{l}}(t,z,\lambda)$*,* $\bar{l} \in \mathcal{Z}_l^n$*, and any* $W(t,z,\lambda) \in \mathfrak{G}^{\,sing}$*, the hypergeometric integral* (*6*) *is a univalued meromorphic function of variables* p, z, λ *holomorphic on the set* (*7*).

For a function $W(t,z,\lambda) \in \mathfrak{G}$, let $\Psi_W(z,\lambda)$ be the following $V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n}$-valued function

$$\Psi_W(z,\lambda) \;=\; \sum_{l_1+\dots+l_n=l} I(w_{\bar{l}}, W)(z,\lambda)\, f^{l_1} v_1 \otimes \dots \otimes f^{l_n} v_n\,. \tag{8}$$

Theorem 9. *(Corollaries 5.25, 5.26 in* [TV1]*.) Let* p, z, λ *be such that conditions* (*7*) *are satisfied.*

i) Let $\mathrm{Im}\,\mu \neq 0$*. Then for any function* $W \in \mathfrak{G}$*, the function* $\Psi_W(z,\lambda)$ *is a solution of the qKZ equation with values in* $(V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n})_l$*.*

ii) Let $\mathrm{Im}\,\mu = 0$*. Then for any function* $W \in \mathfrak{G}^{\,sing}$*, the function* $\Psi_W(z,\lambda)$ *is a solution of the qKZ equation with values in* $(V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n})_l^{sing}$*.*

The solutions of the qKZ equation defined by (8) are called the *hypergeometric solutions.*

4. Main results

Fix natural numbers n, l. We will often assume the following restrictions on the parameters p, z, λ:

$$\mathrm{Re}\,p < 0, \qquad 1 \notin p\mathbb{Z}, \tag{9}$$

$$\{s \mid s \in \mathbb{Z}_{>0}, s < 2\max\{\mathrm{Re}\,\lambda_1,\dots,\mathrm{Re}\,\lambda_n\},\; s \leq l\} \bigcap \{p\mathbb{Z}\} = \emptyset, \tag{10}$$

$$\{2\lambda_m - s \mid s \in \mathbb{Z}_{\geq 0},\; s < 2\mathrm{Re}\,\lambda_m,\; s < l\} \bigcap \{p\mathbb{Z}\} = \emptyset, \qquad m = 1,\dots,n, \tag{11}$$

$$z_k - z_m \pm (\lambda_k + \lambda_m) + s \notin \{p\mathbb{Z}\}, \qquad k,m = 1,\dots,n,\; k \neq m,\; s = 1-l,\dots,l-1. \tag{12}$$

Sometimes, in addition we will make the following assumption. Let $\lambda \in \mathbb{C}^n$. For each $i \in \{1,\dots,n\}$ such that $\lambda_i \notin \Lambda^+$, assume

$$\{1,\dots,l\} \bigcap \{p\mathbb{Z}\} = \emptyset, \tag{13}$$

$$\{2\lambda_i - s \mid s = 0,1,\dots,l-1\} \bigcap \{p\mathbb{Z}\} = \emptyset. \tag{14}$$

Notice that if the parameters p, z, λ satisfy conditions (9)-(14), then they also satisfy conditions (7).

4.1. **Analytic continuation. The case** $\operatorname{Im}\mu \neq 0$**.** Let $\Lambda = (\Lambda_1, \dots, \Lambda_n) \in \mathbb{C}^n$, $\bar{l} = (l_1, \dots, l_n) \in \mathbb{Z}^n_{\geq 0}$. An i-th coordinate of $\bar{l}$ is called Λ-*admissible* if either $\Lambda_i \notin \Lambda^+$ or $\Lambda_i \in \Lambda^+$ and $l_i \leq 2\Lambda_i$. The index $\bar{l}$ is called Λ-*admissible* if all its coordinates are Λ-admissible. Denote $B_\Lambda(\bar{l}) \subset \{1, \dots, n\}$ the set of all non-Λ-admissible coordinates of $\bar{l}$.

Remark. Let p be generic, $\Lambda \in \mathbb{C}^n$, $\bar{l} \in \mathcal{Z}^n_l$. Then the weight coefficient $c_{\bar{l}}(\Lambda)$ is not equal to zero if and only if the index $\bar{l}$ is Λ-admissible.

Let $\bar{m} \in \mathcal{Z}^n_l$ and $B_\Lambda(\bar{l}) \subseteq B_\Lambda(\bar{m})$. Then the function $c(\lambda) = c_{\bar{m}}(\lambda)/c_{\bar{l}}(\lambda)$ is holomorphic at Λ. Moreover, $c(\Lambda) = 0$ if $B_\Lambda(\bar{l}) \neq B_\Lambda(\bar{m})$, and $c(\Lambda)$ is not zero if $B_\Lambda(\bar{l}) = B_\Lambda(\bar{m})$.

Theorem 10. *Let* $\operatorname{Im}\mu \neq 0$*. Let* p *satisfy condition* (9)*. Let* $Z, \Lambda \in \mathbb{C}^n$ *satisfy conditions* (10)-(12)*. Let* $\bar{l}, \bar{m} \in \mathcal{Z}^n_l$*. Assume that for all* $i = 1, \dots, n$ *either the* i*-th coordinate of* $\bar{l}$ *or the* i*-th coordinate of* $\bar{m}$ *is* Λ*-admissible, i.e.* $B(\bar{l}) \bigcap B(\bar{m}) = \emptyset$*. Then the hypergeometric integral* $I(w_{\bar{l}}, W_{\bar{m}})(z, \lambda)$ *is holomorphic at* Z, Λ*. Moreover, there exists a contour of integration* $\mathfrak{I}(Z, \Lambda) \subset \mathbb{C}^l$ *independent on* $\bar{l}, \bar{m}$*, such that for all* z, λ*, in a small neighborhood of* Z, Λ *we have*

$$I(w_{\bar{l}}, W_{\bar{m}})(z, \lambda) = \int\limits_{\mathfrak{I}(Z,\Lambda)} \Phi_l(t, z, \lambda) w_{\bar{l}}(t, z, \lambda) W_{\bar{m}}(t, z, \lambda)\, d^l t.$$

A contour $\mathfrak{I}(Z, \Lambda)$ with properties indicated in Theorem 10 is called an *integration contour associated to* Z, Λ.

Theorem 10 is proved in Section 5.

For $\bar{l}, \bar{m} \in \mathcal{Z}^n_l$, introduce a function

$$J_{\bar{l},\bar{m}}(z, \lambda) = c_{\bar{m}}(\lambda) I(w_{\bar{l}}, W_{\bar{m}})(z, \lambda). \tag{15}$$

Theorem 11. *Let* $\operatorname{Im}\mu \neq 0$*. Let* p *satisfy condition* (9)*. Let* $Z, \Lambda \in \mathbb{C}^n$ *satisfy conditions* (10)-(12)*. Let* $\bar{l}, \bar{m} \in \mathcal{Z}^n_l$*. Then the meromorphic function* $J_{\bar{l},\bar{m}}(z, \lambda)$ *is holomorphic at* Z, Λ*. Moreover, if* $B_\Lambda(\bar{l}) \subset B_\Lambda(\bar{m})$ *and* $B_\Lambda(\bar{l}) \neq B_\Lambda(\bar{m})$*, then* $J_{\bar{l},\bar{m}}(Z, \Lambda) = 0$*.*

Theorem 11 is proved in Section 5.

Let $\Lambda \in \mathbb{C}^n$, $\bar{l}, \bar{m} \in \mathcal{Z}^n_l$ and $B_\Lambda(\bar{l}) = B_\Lambda(\bar{m})$. Denote B the set $B_\Lambda(\bar{l}) = B_\Lambda(\bar{m})$. Introduce $\Lambda'(B) = (\Lambda'_1, \dots, \Lambda'_n) \in \mathbb{C}^n$ by the rule: $\Lambda'_i = \Lambda_i$ if $i \notin B$ and $\Lambda'_i = -\Lambda_i - 1$ if $i \in B$.

Let $\bar{l}'(B) = (l'_1, \dots, l'_n)$, where $l'_i = l_i$ if $i \notin B$ and $l'_i = l_i - 2\Lambda_i - 1$ if $i \in B$. Similarly, let $\bar{m}'(B) = (m'_1, \dots, m'_n)$, where $m'_i = m_i$ if $i \notin B$ and $m'_i = m_i - 2\Lambda_i - 1$ if $i \in B$.

We have $l'_1 + \ldots + l'_n = m'_1 + \ldots + m'_n = l - 2\sum\limits_{i \in B} \Lambda_i - |B|$. We denote this number $l'(B)$.

Theorem 12. *Let* $\operatorname{Im}\mu \neq 0$*. Let* p *satisfy condition* (9)*. Let* $Z, \Lambda \in \mathbb{C}^n$ *satisfy conditions* (10)-(12)*. Let* $\bar{l}, \bar{m} \in \mathcal{Z}^n_l$ *and* $B_\Lambda(\bar{l}) = B_\Lambda(\bar{m}) = B$*. Then*

$$J_{\bar{l},\bar{m}}(Z, \Lambda) = \mathfrak{C}_\Lambda(Z) J_{\bar{l}'(B),\bar{m}'(B)}(Z, \Lambda'(B)),$$

where $\mathfrak{C}_\lambda(z)$ *is a nonzero holomorphic function at* Z *given below.*

Theorem 12 is proved in Section 5.

Notice that $J_{\bar{l}'(B),\bar{m}'(B)}(z, \Lambda'(B)) = c_{\bar{m}'(B)}(\Lambda'(B)) I(w_{\bar{l}'(B)}, W_{\bar{m}'(B)})(z, \Lambda'(B))$, where $I(w_{\bar{l}'(B)}, W_{\bar{m}'(B)})(z, \lambda)$ is an $l'(B)$-dimensional hypergeometric integral. The indices $\bar{l}'(B)$,

$\bar{m}'(B)$ are $\Lambda'(B)$-admissible. By Theorem 10 the integral $I(w_{\bar{l}'(B)}, W_{\bar{m}'(B)})(z, \Lambda'(B))$ is well defined. Theorem 12 connects l- and $l'(B)$-dimensional hypergeometric integrals.

Now, we describe the function $\mathfrak{C}_\lambda(z)$. For $k \in \mathbb{Z}_{\geq 0}$, let

$$\psi_k = \frac{-1}{\pi^{k+2}\, k!\, (k+1)!}\, \Gamma(-(k+1)/p)\, \Gamma(1/p)^{k+1} \prod_{j=1}^{k+1} \sin(j\pi/p).$$

For $z, \lambda \in \mathbb{C}^n, j \in \{1, \dots, n\}, k \in \mathbb{Z}_{\geq 0}$, let

$$\varphi_{\lambda,j,k}(z) = \prod_{s=0}^{k} \left(\prod_{i=0}^{j-1} \frac{\Gamma((z_i - z_j + \lambda_i + \lambda_j - s)/p)}{\Gamma((z_i - z_j - \lambda_i - \lambda_j + s)/p)} \prod_{i=j+1}^{n} \frac{\Gamma((z_j - z_i + \lambda_j + \lambda_i - s)/p)}{\Gamma((z_j - z_i - \lambda_j - \lambda_i + s)/p)} \right).$$

Then

$$\mathfrak{C}_\lambda(z) = \frac{l!}{(l'(B))!} \prod_{j \in B} e^{\mu(2\lambda_j+1) z_j \pi i/p} \psi_{2\lambda_j}\, \varphi_{\lambda,j,2\lambda_j}(z).$$

Consider a square matrix

$$J^l(z, \lambda) = \{J_{\bar{l},\bar{m}}(z, \lambda)\}_{\bar{l},\bar{m} \in \mathcal{Z}_l^n}. \tag{16}$$

Recall that for generic p, we have $c_{\bar{m}}(\lambda) \neq 0$ if $\bar{m}$ is λ-admissible. According to Theorem 7, the matrix (16) is holomorphic if conditions (7) hold. According to Theorem 11, the matrix (16) is holomorphic on a larger set of parameters (9)-(12).

According to Theorem 11, if some of $\lambda_1, \dots \lambda_n$ become nonnegative half-integers, then the matrix (16) becomes upper block triangular in the following sense.

Let $\Lambda \in \mathbb{C}^n$. Divide the set of indices $\mathcal{Z}_l^n$ into subsets labeled by subsets of $\{1, \dots, n\}$. The subset of $\mathcal{Z}_l^n$ corresponding to a subset $B \subset \{1, \dots, n\}$ consists of all $\bar{l} \in \mathcal{Z}_l^n$ such that $B_\Lambda(\bar{l}) = B$. Since the rows and columns of matrix (16) are labeled by elements of $\mathcal{Z}_l^n$, matrix (16) is divided into blocks labeled by pairs $B_1, B_2 \subset \{1, \dots, n\}$.

Choose any order $<$ on the set of all subsets of $\{1, \dots, n\}$ such that for any $B_1, B_2 \subset \{1, \dots, n\}, B_1 \subset B_2$, we have $B_1 < B_2$. Theorem 11 says that matrix (16) is upper block triangular at (z, Λ) with respect to this order. Namely, the block corresponding to a pair B_1, B_2 is equal to zero if $B_1 < B_2$.

Theorem 12 describes the diagonal blocks of matrix (16). The diagonal block of $J^l(z, \Lambda)$ corresponding to a subset $B \subset \{1, \dots, n\}$ coincides (up to multiplication from the left by the non-degenerate diagonal matrix $\{\mathfrak{C}_{\Lambda,\bar{m}}(z)\}_{\bar{m} \in \mathcal{Z}_l^n}$) with the diagonal block of the matrix $J^{l'(B)}(z, \Lambda'(B))$ corresponding to the empty subset of $\{1, \dots, n\}$.

4.2. **Analytic continuation. The case** $\operatorname{Im}\mu = 0$ is treated similarly.

A function $W \in \mathfrak{G}^{\,sing}$ is called Λ-*admissible* if for all non-Λ-admissible $\bar{m} \in \mathcal{Z}_l^n$, the functions $a_{\bar{m}}(\lambda, u)$ in decomposition (1) are equal to zero.

Theorem 13. *Let* $\operatorname{Im}\mu = 0$. *Let* p *satisfy condition* (*9*). *Let* $Z, \Lambda \in \mathbb{C}^n$ *satisfy conditions* (*10*)-(*12*). *Let* $\bar{l} \in \mathcal{Z}_l^n$. *Let* $W \in \mathfrak{G}^{\,sing}$ *be* Λ-*admissible. Then the hypergeometric integral* $I(w_{\bar{l}}, W)(z, \lambda)$ *is holomorphic at* Z, Λ. *Moreover, there exists a contour of integration* $\mathfrak{I}(Z, \Lambda) \subset \mathbb{C}^l$ *independent on* $\bar{l}$ *and* W, *such that for all* z, λ, *in a small neighborhood of* Z, Λ *we have*

$$I(w_{\bar{l}}, W)(z, \lambda) = \int\limits_{\mathfrak{I}(Z,\Lambda)} \Phi_l(t, z, \lambda) w_{\bar{l}}(t, z, \lambda) W(t, z, \lambda)\, d^l t.$$

A contour $\mathfrak{I}(Z,\Lambda)$ with properties indicated in Theorem 13 is called an *integration contour associated to* Z,Λ.

The proof of Theorem 13 is similar to the proof of Theorem 10.

It follows from the proof that a contour of integration associated to Z,Λ with respect to Theorem 10 is also an integration contour associated to Z,Λ with respect to Theorem 13 and vice versa.

Theorem 14. *Let* $\operatorname{Im}\mu=0$. *Let* p *satisfy condition* (*9*). *Let* $Z,\Lambda\in\mathbb{C}^n$ *satisfy conditions* (*10*)-(*12*). *Let* $\bar{l}\in\mathcal{Z}_l^n$, $W\in\mathfrak{G}^{\,sing}$. *Then the meromorphic function* $c_{\bar{l}}(\lambda)I(w_{\bar{l}},W)(z,\lambda)$ *is holomorphic at* Z,Λ. *Moreover, if* $\bar{l}$ *is not* Λ*-admissible and* W *is* Λ*-admissible, then the function* $c_{\bar{l}}(\lambda)I(w_{\bar{l}},W)(z,\lambda)$ *is equal to zero at* (Z,Λ).

The proof of Theorem 14 is similar to the proof of Theorem 11.

Notice that in Theorem 14 we consider the function $c_{\bar{l}}(\lambda)I(w_{\bar{l}},W)(z,\lambda)$ and not the function $c_{\bar{m}}(\lambda)I(w_{\bar{l}},W_{\bar{m}})(z,\lambda)$ as in Theorem 11.

4.3. **Solutions of qKZ with values in irreducible representations.** For $\bar{m}\in\mathcal{Z}_l^n$, consider a $V_{\lambda_1}\otimes\ldots\otimes V_{\lambda_n}$-valued function

$$\Psi_{\bar{m}}(z,\lambda)=\sum_{l_1+\ldots+l_n=l}J_{\bar{l},\bar{m}}(z,\lambda)f^{l_1}v_1\otimes\ldots\otimes f^{l_n}v_n. \tag{17}$$

Corollary 15. *Let* $\operatorname{Im}\mu\neq 0$. *Let* $p\in\mathbb{C}$ *and* $\Lambda\in\mathbb{C}^n$ *satisfy conditions* (*9*)-(*11*). *Then the function* $\Psi_{\bar{m}}(z,\Lambda)$ *given by* (*17*) *is a meromorphic solution of the qKZ equation with values in* $(V_{\Lambda_1}\otimes\ldots\otimes V_{\Lambda_n})_l$, *holomorphic for all* z *satisfying condition* (*12*).

Corollary 15 follows from Theorems 1, 9, and 11.

Let $\Lambda\in\mathbb{C}^n$. For any Λ-admissible $\bar{m}\in\mathcal{Z}_l^n$, consider a function

$$\Psi_{\bar{m}}(z,\Lambda)=\sum I(w_{\bar{l}},W_{\bar{m}})(z,\lambda)f^{l_1}v_1\otimes\ldots\otimes f^{l_n}v_n, \tag{18}$$

where the sum is over all Λ-admissible $\bar{l}\in\mathcal{Z}_l^n$.

Corollary 16. *Let* $\operatorname{Im}\mu\neq 0$. *Let* $p\in\mathbb{C}$ *and* $\Lambda\in\mathbb{C}^n$ *satisfy conditions* (*9*)-(*11*). *Then for any* Λ*-admissible* $\bar{m}\in\mathcal{Z}_l^n$, *the function* $\Psi_{\bar{m}}(z,\Lambda)$ *given by* (*18*) *is a meromorphic solution of the qKZ equation with values in* $(L_{\Lambda_1}\otimes\ldots\otimes L_{\Lambda_n})_l$, *holomorphic for all* z *satisfying condition* (*12*).

Notice that vectors $f^{l_1}v_1\otimes\ldots\otimes f^{l_n}v_n$ with Λ-admissible $\bar{l}\in\mathcal{Z}_l^n$ form a basis in $(L_{\Lambda_1}\otimes\ldots\otimes L_{\Lambda_n})_l$.

Corollary 16 follows from Corollary 15 and Lemma 3.

Let $\operatorname{Im}\mu=0$. For $W\in\mathfrak{G}^{\,sing}$, consider a $V_{\lambda_1}\otimes\ldots\otimes V_{\lambda_n}$-valued function $\Psi_W(z,\lambda)$ given by (8).

Corollary 17. *Let* $\operatorname{Im}\mu=0$. *Let* $p\in\mathbb{C}$ *and* $\Lambda\in\mathbb{C}^n$ *satisfy conditions* (*9*)-(*11*). *Let* $W\in\mathfrak{G}^{\,sing}$ *be* Λ*-admissible. Then the function* $\Psi_W(z,\Lambda)$ *given by* (*8*) *is a meromorpfic solution of the qKZ equation with values in* $(V_{\Lambda_1}\otimes\ldots\otimes V_{\Lambda_n})_l^{sing}$, *holomorphic for all* z *satisfying condition* (*12*).

Corollary 17 follows from Theorems 1, 9, and 13.

Let $\Lambda \in \mathbb{C}^n$, $W \in \mathfrak{G}^{\,sing}$. Consider a function

$$\Psi_W(z,\Lambda) = \sum I(w_{\bar{l}}, W)(z,\lambda) f^{l_1} v_1 \otimes \ldots \otimes f^{l_n} v_n, \tag{19}$$

where the sum is over Λ-admissible $\bar{l} \in \mathcal{Z}_l^n$.

Corollary 18. *Let* $\operatorname{Im}\mu = 0$. *Let* $p \in \mathbb{C}$ *and* $\Lambda \in \mathbb{C}^n$ *satisfy conditions* (*9*)-(*11*). *Let* $W \in \mathfrak{G}^{\,sing}$ *be* Λ*-admissible. Then* $\Psi_W(z,\Lambda)$ *given by* (*19*) *is a meromorphic solution of the qKZ equation with values in* $(L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})_l^{sing}$, *holomorphic for all* z *satisfying condition* (*12*).

Corollary 18 follows from Corollary 17 and Lemma 3.

The solutions of the qKZ equation defined by formulas (18),(19) are called the *hypergeometric solutions.*

4.4. **Determinant of the hypergeometric pairing.** Let $\operatorname{Im}\mu \neq 0$. The determinant of the matrix $J^l(z,\lambda)$ defined by (16) is given by

$$\begin{aligned}
\det(J_l(z,\lambda)) = (2i)^{l\binom{n+l-1}{n-1}} (l!)^{\binom{n+l-1}{n-1}} (e^{\mu}-1)^{-2\sum_{m=1}^{n} \lambda_m/p\cdot\binom{n+l-1}{n} + 2n/p\cdot\binom{n+l-1}{n+1}} \times \\
\times \exp(\mu \sum_{m=1}^{n} z_m/p \cdot \binom{n+l-1}{n} \times \\
\times \exp\left((\mu+\pi i) \left(\sum_{m=1}^{n} \lambda_m/p \cdot \binom{n+l-1}{n} - n/p \cdot \binom{n+l-1}{n+1} \right) \right) \times \\
\times \prod_{s=0}^{l-1} \Bigg[\Gamma(1+1/p)^n \Gamma(1+(s+1)/p)^{-n} \prod_{m=1}^{n} \frac{\pi}{\Gamma(1-(2\lambda_m - s)/p)} \times \\
\times \prod_{1\le k<m\le n} \frac{\Gamma((z_k+\lambda_k-z_m+\lambda_m-s)/p)}{\Gamma((z_k-\lambda_k-z_m-\lambda_m+s)/p)} \Bigg]^{\binom{n+l-s-2}{n-1}} ,
\end{aligned}$$

cf. Theorem 5.14 of [TV1].

For $\Lambda \in \mathbb{C}^n$, consider a matrix

$$J^l_{\mathrm{adm}}(z,\lambda) = \{ J_{\bar{l},\bar{m}}(z,\lambda) \}, \tag{20}$$

where $\bar{l}, \bar{m} \in \mathcal{Z}_l^n$ run through the set of all Λ-admissible indices.

By Theorem 10, the matrix $J^l_{\mathrm{adm}}(z,\lambda) = J^l_{\mathrm{adm}}(z,\lambda,p)$ is holomorphic at (z,Λ) if parameters p, z, Λ satisfy conditions (9)-(12).

Theorem 19. *Let* $\operatorname{Im}\mu \neq 0$. *Let* $p \in \mathbb{C}$ *and* $z, \Lambda \in \mathbb{C}^n$ *satisfy conditions* (*9*)-(*14*). *Then the matrix* $J^l_{\mathrm{adm}}(z,\Lambda)$ *is non-degenerate. Moreover, for generic* p,

$$\det(J^l_{\mathrm{adm}}(z,\Lambda)) = \prod_{A,\, A\subset B(\Lambda)} \Big(C_\Lambda(A)(z) \det(J^{l'(A)}(z,\Lambda'(A))) \Big)^{(-1)^{|A|}} , \tag{21}$$

where $B(\Lambda)$ is the set of all $i \in \{1, \dots, n\}$ such that $\Lambda_i \in \Lambda^+$, and the function $C_\Lambda(A)(z)$ is given by

$$C_\Lambda(A)(z) = \prod_{\bar m,\, B(\bar m) \supseteq A} \; \prod_{i \in A} \left(\frac{l!}{(l - 2\Lambda_i - 1)!} \; \psi_{2\Lambda_i} \, \varphi_{\Lambda, i, 2\Lambda_i}(z) \right).$$

Theorem 19 is proved in Section 5.

Consider a pairing

$$s_\mu(z) : \; (L^q_{\Lambda_1} \otimes \dots \otimes L^q_{\Lambda_n})_l \otimes ((L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n})_l)^* \to \mathbb{C},$$

defined by

$$(f^{m_1} v_1^q \otimes \dots \otimes f^{m_n} v_n^q) \otimes (f^{l_1} v_1 \otimes \dots \otimes f^{l_n} v_n)^* \; \mapsto \; J_{\bar l, \bar m}(z, \Lambda) \in \mathbb{C},$$

where $\bar l, \bar m \in \mathcal{Z}_l^n$ are Λ-admissible.

Corollary 20. *Let $\operatorname{Im} \mu \neq 0$. Let $p \in \mathbb{C}$ and $z, \Lambda \in \mathbb{C}^n$ satisfy conditions (9)-(14). Then the pairing $s_\mu(z)$ is well defined and non-degenerate.*

Corollary 20 follows from Theorem 19.

Corollary 21. *Let $\operatorname{Im} \mu \neq 0$. Let $p \in \mathbb{C}$ and $\Lambda \in \mathbb{C}^n$ satisfy conditions (9)-(11) and (13)-(14). Then any solution of the qKZ equation with values in $(L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n})_l$ is a linear combination of the hypergeometric solutions (18) with p-periodic coefficients.*

Corollary 21 follows from Corollary 20.

Similarly, for a permutation $\sigma \in \mathbb{S}^n$, consider a pairing

$$s^\sigma_\mu(z) : \; (L^q_{\Lambda_{\sigma_1}} \otimes \dots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes ((L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n})_l)^* \to \mathbb{C},$$

defined by

$$(f^{m_{\sigma_1}} v^q_{\sigma_1} \otimes \dots \otimes f^{m_{\sigma_n}} v^q_{\sigma_n}) \otimes (f^{l_1} v_1 \otimes \dots \otimes f^{l_n} v_n)^* \; \mapsto \; c_{\bar m}(\Lambda) I(w_{\bar l}, W^\sigma_{\bar m})(z, \Lambda) \in \mathbb{C},$$

where $\bar l, \bar m \in \mathcal{Z}_l^n$ are Λ-admissible.

Corollary 22. *Let $\operatorname{Im} \mu \neq 0$. Let $p \in \mathbb{C}$ and $z, \Lambda \in \mathbb{C}^n$ satisfy conditions (9)-(14). Then for any permutation $\sigma \in \mathbb{S}^n$, the pairing $s^\sigma_\mu(z)$ is well defined and non-degenerate.*

Corollary 22 follows from Corollaries 20 and 6.

For a permutation $\sigma \in \mathbb{S}^n$ and a Λ-admissible index $\bar m \in \mathcal{Z}_l^n$, consider a function

$$\Psi^\sigma_{\bar m}(z, \Lambda) = \sum I(w_{\bar l}, W^\sigma_{\bar m})(z, \lambda) f^{l_1} v_1 \otimes \dots \otimes f^{l_n} v_n, \tag{22}$$

where the sum is over all Λ-admissible $\bar l \in \mathcal{Z}_l^n$.

Corollary 23. *Let $\operatorname{Im} \mu \neq 0$. Let $\sigma \in \mathbb{S}^n$ be a permutation. Let $p \in \mathbb{C}$ and $\Lambda \in \mathbb{C}^n$ satisfy conditions (9)-(11) and (13)-(14). Then any solution of the qKZ equation with values in $(L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n})_l$ is a linear combination of the hypergeometric solutions (22) with p-periodic coefficients.*

Corollary 23 follows from Corollary 22.

The map $s^\sigma_\mu(z)$ induces a map

$$\tilde s^\sigma_\mu(z) : \; (L^q_{\Lambda_{\sigma_1}} \otimes \dots \otimes L^q_{\Lambda_{\sigma_n}})_l \to (L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n})_l. \tag{23}$$

This map is an isomorphism of vector spaces for all z satisfying (12). For a fixed vector $v^q \in (L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l$, the $(L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})_l$-valued function $\tilde{s}^{\sigma}_{\mu}(z)v^q$ is a solution of the qKZ equation. This construction identifies the space $\mathcal{S}$ of all meromorphic solutions to the qKZ equation with the space $(L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F$ where F is the field of scalar meromorphic functions in $z_1, \ldots, z_n$, p-periodic with respect to each of the variables,

$$\iota_\sigma : \ (L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F \to \mathcal{S}.$$

Let now $\operatorname{Im} \mu = 0$. Consider a pairing

$$s_\mu(z) : \ (L^q_{\Lambda_1} \otimes \ldots \otimes L^q_{\Lambda_n})^{sing}_l \otimes ((L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})_l)^* \to \mathbb{C},$$

defined by

$$v^q \otimes (f^{l_1} v_1 \otimes \ldots \otimes f^{l_n} v_n)^* \ \mapsto \ I(w_{\bar{l}}, \mathfrak{B}_{\mathrm{id}}(z,\Lambda) v^q)(z,\Lambda) \in \mathbb{C},$$

where $\bar{l} \in \mathcal{Z}^n_l$ is admissible, $v^q \in (L^q_{\Lambda_1} \otimes \ldots \otimes L^q_{\Lambda_n})^{sing}_l$. The map $\mathfrak{B}_{\mathrm{id}}(z,\Lambda)$ is defined in (3), the integral $I(w_{\bar{l}}, \mathfrak{B}_{\mathrm{id}}(z,\Lambda)v^q)(z,\Lambda)$ is taken over an integration contour associated to (z,Λ).

The map $s_\mu(z)$ induces a map

$$\tilde{s}_\mu(z) : \ (L^q_{\Lambda_1} \otimes \ldots \otimes L^q_{\Lambda_n})^{sing}_l \to (L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})_l. \tag{24}$$

By Corollary 18, the image of $\tilde{s}_\mu(z)$ belongs to $(L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})^{sing}_l$.

Theorem 24. *Let* $\operatorname{Im} \mu = 0$. *Let* $p \in \mathbb{C}$ *and* $z, \Lambda \in \mathbb{C}^n$ *satisfy conditions (9)-(14). Let* $\sum_{m=1}^{n} 2\Lambda_m - s \notin \{p\mathbb{Z}_{<0}\}$ *for* $s = l-1, \ldots, 2l-2$. *Then the map*

$$\tilde{s}_\mu(z) : \ (L^q_{\Lambda_1} \otimes \ldots \otimes L^q_{\Lambda_n})^{sing}_l \to (L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})^{sing}_l$$

is an isomorphism of vector spaces.

Theorem 24 is proved similarly to Theorem 19 using Theorem 5.15 in [TV1].

Corollary 25. *Let* $\operatorname{Im} \mu = 0$. *Let* $p \in \mathbb{C}$ *and* $\Lambda \in \mathbb{C}^n$ *satisfy conditions (9)-(11) and (13)-(14). Let* $\sum_{m=1}^{n} 2\Lambda_m - s \notin \{p\mathbb{Z}_{<0}\}$ *for* $s = l-1, \ldots, 2l-2$. *Then any solution of the qKZ equation with values in* $(L_{\Lambda_1} \otimes \ldots \otimes L_{\Lambda_n})^{sing}_l$ *is a linear combination of the hypergeometric solutions (19) with p-periodic coefficients.*

Corollary 25 follows from Theorem 24.

4.5. **Asymptotic solutions.** Let U be a domain in $\mathbb{C}^n$ and let $M_1, \ldots, M_n$ be sl_2 modules. A basis $\Psi_1, \ldots, \Psi_N$ of solutions to the qKZ equation with values in $(M_1 \otimes \ldots \otimes M_n)_l$, is called an *asymptotic solution* in the domain U if

$$\Psi_j(z) = \exp\Big(\sum_{m=1}^{n} a_{mj} z_m / p\Big) \prod_{1 \le m < k \le n} (z_k - z_m)^{b_{jkm}} (v_j + o(1)),$$

where a_{mj} and b_{jkm} are suitable numbers, $v_1, \ldots, v_N$ are vectors which form a basis in $(M_1 \otimes \ldots \otimes M_n)_l$ and $o(1)$ tends to zero as z tends to infinity in U. The domain U is called an *asymptotic zone.*

Let $\Psi^1_1(z),\dots,\Psi^1_N(z)$ and $\Psi^2_1(z),\dots,\Psi^2_N(z)$ be asymptotic solutions in asymptotic zones $U_1 \in \mathbb{C}^n$ and $U_2 \in \mathbb{C}^n$, respectively. Then

$$\Psi^1_i(z) \;=\; \sum_{j=1}^{N} T^j_i(z)\,\Psi^2_j(z), \qquad i=1,...,N$$

for suitable meromorphic functions $T^j_i(z)$. The matrix valued function $T(z)=(T^j_i(z))$ is p periodic with respect to variables $z_1,...,z_n$ and has nonzero determinant for generic z. The function $T(z)$ is called *the transition function between the asymptotic solutions* Ψ^1 *and* Ψ^2.

We describe the asymptotic zones, asymptotic solutions and transition functions of the qKZ equation with values in $(L_{\lambda_1}\otimes\dots\otimes L_{\lambda_n})_l$. Our results are parallel to the results of [TV1], where the asymptotic zones, asymptotic solutions and transition functions are described for the qKZ equation with values in $(V_{\lambda_1}\otimes\dots\otimes V_{\lambda_n})_l$.

For a permutation $\sigma\in\mathbb{S}^n$, define an asymptotic zone in $\mathbb{C}^n$,

$$U_\sigma=\{z\in\mathbb{C}^n\,|\,\mathrm{Re}\,z_{\sigma_1}\ll\dots\ll\mathrm{Re}\,z_{\sigma_n}\}. \tag{25}$$

Say that $z\to\infty$ in U_σ if $\mathrm{Re}\,(z_{\sigma_m}-z_{\sigma_{m+1}})\to-\infty$ for all $m=1,\dots,n-1$.

For each permutation $\sigma\in\mathbb{S}^n$, let $\Psi^\sigma(\lambda)$ be the set of solutions $\{\Psi^\sigma_{\bar m}\}$ of the qKZ equation with values in $(L_{\lambda_1}\otimes\dots\otimes L_{\lambda_n})_l$ given by (22). Here $\bar m\in\mathcal{Z}^n_l$ runs through the set of all λ-admissible indices.

Theorem 26. *Let* $\mathrm{Im}\,\mu\neq 0$. *Let* $p\in\mathbb{C},\Lambda\in\mathbb{C}^n$ *satisfy the conditions* (*9*)-(*11*). *Then for any permutation* $\sigma\in\mathbb{S}^n$, *the set of solutions* $\Psi^\sigma(\Lambda)$ *forms a basis of solutions. Moreover, the basis of solutions* $\Psi^\sigma(\Lambda)$ *is an asymptotic solution of the qKZ equation with values in* $(L_{\Lambda_1}\otimes\dots\otimes L_{\Lambda_n})_l$ *in the asymptotic zone* U_σ. *Namely,*

$$\Psi^\sigma_{\bar l}(z)=\Xi_{\bar l}\,\exp(\mu\sum_{m=1}^{n}l_m z_m/p)\times$$
$$\times\prod_{1\le k<m\le n}((z_{\sigma_k}-z_{\sigma_m})/p)^{2(l_{\sigma_k}\Lambda_{\sigma_m}+l_{\sigma_m}\Lambda_{\sigma_k}-l_{\sigma_k}l_{\sigma_m})/p}(f^{l_1}v_1\otimes\dots\otimes f^{l_n}v_n+o(1))$$

as $z\to\infty$ *in* U_σ *so that at any moment condition* (*12*) *is satisfied. Here* $|\arg((z_{\sigma_k}-z_{\sigma m})/p)|<\pi$ *and* $\Xi_{\bar l}$ *is a constant independent of the permutation* σ *and given by*

$$\Xi_{\bar l}=(2i)^l\,l!\,\Gamma(-1/p)^{-l}\prod_{m=1}^{n}\big[(e^\mu-1)^{(l_m(l_m-1)-2l_m\Lambda_m)/p}\times$$
$$\times\exp((\mu+\pi i)(l_m\Lambda_m-l_m(l_m-1)/2)/p)\prod_{s=1}^{l_m-1}\Gamma((2\Lambda_m-s)/p)\Gamma(-(s+1)/p)\big],$$

where $0\le\arg(e^\mu-1)<2\pi$.

The proof of Theorem 26 is the same as the proof of Theorem 6.4 in [TV1].

Notice that the asymptotics of the basis of solutions $\Psi^\sigma(\Lambda)$ determine the basis uniquely. Namely, if a basis of solutions meromorphically depends on parameters μ,z,λ and has asymptotics in U_σ described in Theorem 26, then such a basis coincides with the basis $\Psi^\sigma(\Lambda)$, see the Remark after Theorem 6.4 in [TV1].

The isomorphism $\iota_\sigma : (L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F \to \mathcal{S}$, defined in Section 4.4, allows us to identify the basis of solutions $\Psi^\sigma(\Lambda)$ with a basis of $(L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F$, considered as a vector space over F. Then the transition function between two asymptotic solutions $\Psi^\sigma(\Lambda)$ and $\Psi^\nu(\Lambda)$ is identified with an F-linear map $(L^q_{\Lambda_{\nu_1}} \otimes \ldots \otimes L^q_{\Lambda_{\nu_n}})_l \otimes F \to (L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F$.

Corollary 27. *Let* $\operatorname{Im}\mu \neq 0$. *Let* $p \in \mathbb{C}, \Lambda \in \mathbb{C}^n$ *satisfy the conditions* (*9*)-(*12*) *and* (*13*)-(*14*). *Then for any permutation* $\sigma \in \mathbb{S}^n$ *and a simple transposition* $(m, m+1)$, *the transition function between asymptotic solutions* $\Psi^\sigma(\Lambda)$ *and* $\Psi^{\sigma\circ(m,m+1)}(\Lambda)$,

$$(L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_{m+1}}} \otimes L^q_{\Lambda_{\sigma_m}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F \to (L^q_{\Lambda_{\sigma_1}} \otimes \ldots \otimes L^q_{\Lambda_{\sigma_n}})_l \otimes F,$$

equals the operator $P_{L^q_{\lambda_{\sigma_{m+1}}} L^q_{\lambda_{\sigma_m}}} R_{L^q_{\lambda_{\sigma_{m+1}}} L^q_{\lambda_{\sigma_m}}}(e^{2\pi i(z_{\sigma_{m+1}} - z_{\sigma_m})/p})$ *acting in the* m*-th and* $(m+1)$*-th factors.*

Corollary 27 follows from Theorem 26 and Corollary 6.

Theorem 26 allows us to obtain another formula for the determinant of the hypergeometric pairing (cf. formula (21)).

Theorem 28. *(Joint with V.Tarasov) Let* $\operatorname{Im}\mu \neq 0$. *Let* $p \in \mathbb{C}$ *and* $z, \Lambda \in \mathbb{C}^n$ *satisfy conditions* (*9*)-(*14*). *Then the matrix* $J^l_{\mathrm{adm}}(z, \Lambda)$ *defined in* (*20*) *is non-degenerate. Moreover, for generic* p,

$$\det(J^l_{\mathrm{adm}}(z,\Lambda)) =$$
$$\mathcal{D}_l(\Lambda)\exp(\mu\sum_{m=1}^n D_m(\Lambda) z_m/p) \prod_{1\le k<m\le n} \prod_{s=0}^{d_{km}} \left(\frac{\Gamma((z_k+\lambda_k-z_m+\lambda_m-s)/p)}{\Gamma((z_k-\lambda_k-z_m-\lambda_m-s)/p)}\right)^{E_{km}(s,\Lambda)},$$

where $d_{km} = \min\{l, \dim L_{\Lambda_k} - 1, \dim L_{\Lambda_m} - 1\}$, *the functions* $\mathcal{D}_l(\Lambda)$, $D_m(\Lambda)$ *and* $E_{km}(s,\Lambda)$ *are given below.*

For $m = 1, \ldots, n$ and $\Lambda \in \mathbb{C}^n$, set

$$D_m(\Lambda) = \sum_{r=1}^{\min\{l, \dim L_{\Lambda_m} - 1\}} (r \dim(L_{\Lambda_1} \otimes \ldots \otimes \widehat{L_{\Lambda_m}} \otimes \ldots \otimes L_{\Lambda_n})_{l-r}).$$

For $k, m = 1, \ldots, n$, $s \in \mathbb{Z}_{\ge 0}$ and $\Lambda \in \mathbb{C}^n$, set

$$E_{km}(s,\Lambda) = \sum_r (\dim(L_{\Lambda_k} \otimes L_{\Lambda_m})_r - s - 1) \dim(L_{\Lambda_1} \otimes \ldots \otimes \widehat{L_{\Lambda_k}} \otimes \ldots \otimes \widehat{L_{\Lambda_m}} \otimes \ldots \otimes L_{\Lambda_n})_{l-r},$$

where the sum is over $r = s+1, \ldots, \min\{l, \dim L_{\Lambda_k} + \dim L_{\Lambda_k} - 1 - s\}$.

Finally, for $\Lambda \in \mathbb{C}^n$, set

$$\mathcal{D}_l(\Lambda) = \prod_{\bar{l}} c_{\bar{l}}\, \Xi_{\bar{l}},$$

where the product is over Λ-admissible indices $\bar{l} \in \mathcal{Z}^n_l$.

The proof of Theorem 28 is the same as the proof of Theorem 6.4 in [TV1].

5. Proofs.

In this Section we collected the proofs of statements from Section 4.

5.1. **Proof of Theorem 10.** Let $\operatorname{Im}\mu \neq 0$. Fix a rational weight function $w = w_{\bar l}$, and a trigonometric weight function $W = W_{\bar m} \in \mathfrak{G}$, such that $B(\bar l) \bigcap B(\bar m) = \emptyset$.

For $z, \lambda \in \mathbb{C}^n$ such that $\operatorname{Re}(z_i + \lambda_i) < 0$ and $\operatorname{Re}(z_i - \lambda_i) > 0$ for all $i = 1, \dots, n$, the integral $I(w, W)(z, \lambda)$ is defined by formula (4),

$$I(w,W)(z,\lambda) = \int\limits_{\substack{\operatorname{Re} t_i = 0,\\ i=1,\dots,l}} f(t,z,\lambda)\, d^l t,$$

where $f(t,z,\lambda) = \Phi_l(t,z,\lambda) w(t,z,\lambda) W(t,z,\lambda)$. The integrand f has simple poles at

$$t_i = z_j \pm (\lambda_j + kp), \qquad k \in \mathbb{Z}_{\geq 0}, \qquad i = 1, \dots, l, \qquad j = 1, \dots, n, \tag{26}$$

and at

$$t_i - t_j = \pm(1 - kp), \qquad k \in \mathbb{Z}_{\geq 0}, \qquad i, j = 1, \dots, l, \qquad i < j. \tag{27}$$

On the complex line the poles of the first type and the integration contour can be represented as follows.



The integral $I(w,W)(z,\lambda) = I(w,W)(z,\lambda,p)$ is a meromorpic function of z, λ, p.

Fix $Z, \Lambda \in \mathbb{C}^n$, $P \in \mathbb{C}$, satisfying conditions (9)-(12). Our goal is to prove that $I(w,W)(Z,\Lambda,P)$ is well defined and is given by the integral over a suitable cycle.

We fix parameters z^0, λ^0, p^0 in such a way that $\operatorname{Re}\lambda_k^0 < 0$, p^0 is a negative number with large absolute value and $|\operatorname{Im}(z_k^0 + \Lambda_k - z_m^0 - \Lambda_m)|$ is large for $k, m = 1, \dots, n$ and $k \neq m$. Namely, let $p^0 = -2\sum\limits_{j=1}^{n} |\Lambda_j| - 1$, $z_k^0 = i\,(-\operatorname{Im}\Lambda_k + 3kA)$, $\lambda_k^0 = -1 + i\operatorname{Im}\Lambda_k$, and A is a large real number such that $A > 2|\Lambda_k|$, $k = 1, \dots, n$.

The proof is done according to the following plan. First, we represent the integral $I(w,W)(z^0,\lambda^0,p^0)$ as a sum of new integrals of different dimensions. Then we analytically continue the function $I(w,W)(z,\lambda,p)$ from z^0, λ^0, p^0 to z^0, Λ, p^0. Then we analytically continue the function $I(w,W)(z,\Lambda,p)$ from z^0, Λ, p^0 to z^0, Λ, P. In the last step of the proof we analytically continue the function $I(w,W)(z,\Lambda,P)$ from z^0, Λ, P to Z, Λ, P.

For $\bar k \in \mathbb{Z}^n_{\geq 0}$, set $k = k_1 + \dots + k_n$ and define the multiple residues of f by the formula

$$\begin{aligned}\operatorname{res}_{\bar k} f = {} & \operatorname{res}_{t_k = z_n + \lambda_n - k_n + 1} \dots \operatorname{res}_{t_{k_1+k_2} = z_2 + \lambda_2 - k_2 + 1} \dots \operatorname{res}_{t_{k_1+1} = z_2 + \lambda_2} \\ & \operatorname{res}_{t_{k_1} = z_1 + \lambda_1 - k_1 + 1} \dots \operatorname{res}_{t_2 = z_1 + \lambda_1 - 1} \operatorname{res}_{t_1 = z_1 + \lambda_1} f.\end{aligned} \tag{28}$$

If $k > l$, then we set $\operatorname{res}_{\bar k} f = 0$.

The function $\operatorname{res}_{\bar k} f$ is a function of variables $t_{k+1}, \dots, t_l; z, \lambda$. In particular, for $\bar k = (0, \dots, 0)$, we have $\operatorname{res}_{\bar k} f = f$.

Lemma 29. *Let $\bar k, \bar k' \in \mathbb{Z}^n_{\geq 0}$. If $\bar k'$ can be obtained by a permutation of components of $\bar k$ then $\operatorname{res}_{\bar k} f = \operatorname{res}_{\bar k'} f$.*

Lemma 29 follows from the symmetry of the integrand f, see Remark in Section 3.2.

Lemma 30. *For all $i, j = 1, \dots, l$, and for any $a \in \mathbb{C}$, we have*

$$\mathrm{res}_{t_j=a}\, \mathrm{res}_{t_i=a}\, f = 0.$$

Proof: Recall that $f = \Phi_l w W$. On the hyperplane $t_i = t_j$, any function which is symmetric with respect to either rational or trigonometric action is identically equal to zero. In particular, w is equal to zero at the hyperplane $t_i = t_j$. Since all poles of f are at most of first order, we have the statement of the Lemma. □

Lemma 31. *Let $\bar{k} \in \mathbb{Z}^n_{\geq 0}$. The function $\mathrm{res}_{\bar{k}} f$ is holomorpic for all $t_{k+1}, \dots, t_l; z, \lambda, p$ satisfying conditions (9) and such that*

$$t_i \neq z_j \pm (\lambda_j + sp), \qquad s \in \mathbb{Z}_{\geq 0}, \qquad i = k+1, \dots, l, \qquad j = 1, \dots, n,$$

$$t_i - t_j \neq \pm(1 - sp), \qquad s \in \mathbb{Z}_{\geq 0}, \qquad i, j = k+1, \dots, l, \qquad i < j,$$

$$t_i \neq z_j + \lambda_j - m \pm (1 - sp), \quad s \in \mathbb{Z}_{\geq 0}, \quad i = k+1, \dots, l, \quad j = 1, \dots, n, \quad m = 0, \dots, k_j,$$

$$z_i + \lambda_i - z_j \pm (\lambda_j + sp) \neq m, \qquad s \in \mathbb{Z}_{\geq 0}, \qquad i, j = 1, \dots, k, \qquad m = 0, \dots, k, \qquad i \neq j,$$

$$2\lambda_i + sp \neq m, \qquad s \in \mathbb{Z}_{\geq 0}, \qquad i = 1, \dots, k, \qquad m = 0, \dots, k_i.$$

Lemma 31 follows from formulas (26),(27) for the poles of f, and formula (28) for the function $\mathrm{res}_{\bar{k}} f$.

For $u \in \mathbb{R}^n$, define a curve $\mathcal{C}_u = \{\mathcal{C}_u(x) \in \mathbb{C} \,|\, x \in \mathbb{R}\}$ with the following properties. The curve $\mathcal{C}_u$ consists of $2n+1$ line segments and none of the line segments is horizontal (i.e. no line-segment is parallel to the real line). There are two line segments (rays) which go to $+i\infty$ and $-i\infty$, these two rays are parts of the imaginary line. There are n vertical segments with real coordinates $u_1, \dots, u_n$ and imaginary coordinates close to $\mathrm{Im}\,(z^0_1 + \lambda^0_1), \dots, \mathrm{Im}\,(z^0_n + \lambda^0_n)$.

The precise formulas defining $\mathcal{C}_u$ are as follows. For $x \leq A$ and for $x \geq (3n-1)A$, let $\mathcal{C}_u(x) = ix$. For $(3m-1)A \leq x \leq (3m+1)A$, $m = 1, \dots, n$, let $\mathcal{C}_u(x) = u_m + ix$, and this segment is called the *m-th vertical segment.* For $m = 0, \dots, n$, let $\{\mathcal{C}_u(x), (3m+1)A \leq x \leq (3m+2)A\}$ be a parametrization of the line segment connecting $\mathcal{C}_u((3m+1)A)$ and $\mathcal{C}((3m+2)A)$.

See an example of such a curve on the Figure of the proof of Lemma 33.

Lemma 32. *Let $, \Lambda, z^0, \lambda^0, p^0$ be as above. For any sufficiently small ϵ,*

$$I(w, W)(z^0, \lambda^0, p^0) = \sum_{\bar{k}} \frac{l!}{n!\, k_1! \dots k_n! (l-k)!} \int\limits_{\substack{t_j \in \mathcal{C}, \\ j = k+1, \dots, l}} \mathrm{res}_{\bar{k}} f(t, z^0, \lambda^0, p^0)\, d^{l-k} t, \qquad (29)$$

where $d^{l-k} t = dt_{k+1} \dots dt_l$. Here $\mathcal{C}$ denotes the curve $\mathcal{C}_u$ with $u_m = \min\{-\mathrm{Re}\,(2\Lambda_m - \lambda^0_m) - \epsilon, 0\}$, $m = 1, \dots, n$, and the sum is over all $\bar{k} \in \mathbb{Z}^n_{\geq 0}$ such that $k \leq l$, and $k_m \leq 2\mathrm{Re}\,\Lambda_m + 1$ for $m = 1, \dots, n$.

Proof: The integrand decays exponentially with respect to all $t_1, \dots, t_l$. We move the contour of integration with respect to each variable t_i to the left avoiding the poles at $t_i - t_j = (1 - kp)$.

Initially, each of the variables runs through the imaginary line. We deform the contours of integration with respect to $t_1, \dots, t_l$ in such a way that at every moment the contours of integration with respect to $t_1, \dots, t_l$ are curves $\mathcal{C}_{u^i}$ with different parameters $u^i \in \mathbb{R}^n$.

First, we separate the contours of integration with respect to different variables by small distances as follows.

$$I(w,W)(z^0,\lambda^0,p^0) = \int\limits_{\substack{\operatorname{Re} t_j = 0,\\ j=i,\dots,l}} f d^l t = \int\limits_{\substack{t_j \in \mathcal{C}_{u^j},\\ j=1,\dots,l}} f d^l t,$$

where $u^j = ((j-l-1)\rho, \dots, (j-l-1)\rho) \in \mathbb{R}^n$, and ρ is a small positive number.

We deform the integration cycle $\{t \in \mathbb{C}^l \,|\, t_j \in \mathcal{C}_{u^j},\ j = 1, \dots, l\}$ changing u^j, $j = 1, \dots, l$. Namely, we move the vertical segments of contours of integrations $\mathcal{C}_{u^j}$, $j = 1, \dots, n$, to the left, until we reach the vertical segments of the curve $\mathcal{C}$, described in the Lemma, and get the integration cycle $\{t \in \mathbb{C}^l \,|\, t_j \in \mathcal{C},\ j = 1, \dots, l\}$. During the deformation, we keep the same small distances between the contours of integrations $\mathcal{C}_{u^j}$ with respect to different variables. At every moment, when one of the vertical segments of the contour of integration goes through a pole of the integrand, we add another integral of lower dimension which is the integral with respect to remaining variables of the residue. For example, the first such event leads to the following decomposition.

$$\int\limits_{\substack{t_j \in \mathcal{C}_{u^j},\\ j=1,\dots,l}} f d^l t = \int\limits_{\substack{t_j \in \mathcal{C}_{(\operatorname{Re}\lambda_1^0 + j\rho, u_2^j, \dots, u_n^j)},\\ j=1,\dots,l}} f d^l t =$$

$$\int\limits_{\substack{t_1 \in \mathcal{C}_{(\operatorname{Re}\lambda_1^0 - \rho/2, u_2^1, \dots, u_n^1)}\\ t_j \in \mathcal{C}_{(\operatorname{Re}\lambda_1^0 + (j-3/2)\rho, u_2^j, \dots, u_n^j)},\\ j=2,\dots,l}} f d^l t + \int\limits_{\substack{t_j \in \mathcal{C}_{(\operatorname{Re}\lambda_1^0 + (j-3/2)\rho, u_2^j, \dots, u_n^j)},\\ j=2,\dots,l}} \operatorname{res}_{t_1 = z_1 + \lambda_1} f d^{l-1} t.$$

Recall that $\operatorname{Re}\lambda_1^0 = -1$. In the first term of the resulting decomposition, moving the first vertical segment of the contour $\mathcal{C}_{(\operatorname{Re}\lambda_1^0 + 1/2\rho, u_2^2, \dots, u_n^2)}$ corresponding to t_2, through the point $z_1^0 + \lambda_1^0$, we also get a lower dimensional integral. This integral is equal to the second term of the above decomposition by Lemma 29. Note that due to Lemma 30, the integrand of the second term of the decomposition does not have poles at $t_j = z_1 + \lambda_1$ for all $j = 2, \dots, l$. Thus, we can move the first vertical segments of the contours corresponding to t_j, $j = 2, \dots, l$, through $z_1^0 + \lambda_1^0$ without getting new residues.

Finally, we have

$$I(w,W)(z^0,\lambda^0,p^0) =$$
$$\int\limits_{\substack{t_j \in \mathcal{C}_{(\operatorname{Re}\lambda_1^0 + (j-l-1)\rho, u_2^j, \dots, u_n^j)},\\ j=1,\dots,l}} f d^l t + l \int\limits_{\substack{t_j \in \mathcal{C}_{(\operatorname{Re}\lambda_1^0 + (j-l-1)\rho, u_2^j, \dots, u_n^j)},\\ j=2,\dots,l}} \operatorname{res}_{t_1 = z_1 + \lambda_1} f d^{l-1} t.$$

In the first summand we move the contour of integration to

$$\{t \in \mathbb{C}^l \,|\, t_j \in \mathcal{C}_{(\min\{-\mathrm{Re}\,(2\Lambda_1-\lambda_1^0)-\epsilon,0\},u_2^j,\dots,u_n^j)},\ j=1,\dots,l\}$$

and the contour does not meet any other poles of the integrand.

The integrand of the second summand has a simple pole at $t_j = z_1+\lambda_1-1$, since f had a pole at $t_1-t_j=1$. It also may have a pole at $2\lambda_1=0$ coming from the pole at $t_1 = z_1-\lambda_1$, see Lemma 31. So, as before, we have

$$\int\limits_{\substack{t_j\in\mathcal{C}_{(\mathrm{Re}\,\lambda_1^0+(j-l-1)\rho,u_2^j,\dots,u_n^j)},\\ j=2,\dots,l}} \mathrm{res}_{t_1=z_1+\lambda_1} f\, d^{l-1}t = \int\limits_{\substack{t_j\in\mathcal{C}_{(\mathrm{Re}\,\lambda_1^0-1+(j-1)\rho,u_2^j,\dots,u_n^j)},\\ j=2,\dots,l}} \mathrm{res}_{t_1=z_1+\lambda_1} f\, d^{l-1}t =$$

$$\int\limits_{\substack{t_j\in\mathcal{C}_{(\mathrm{Re}\,\lambda_1^0-1+(j-l-1)\rho,u_2^j,\dots,u_n^j)},\\ j=2,\dots,l}} \mathrm{res}_{t_1=z_1+\lambda_1} f\, d^{l-1}t +$$

$$+(l-1) \int\limits_{\substack{t_j\in\mathcal{C}_{(\mathrm{Re}\,\lambda_1^0-1+(j-l-1))\rho,u_2^j,\dots,u_n^j)},\\ j=3,\dots,l}} \mathrm{res}_{t_2=z_1+\lambda_1-1}\mathrm{res}_{t_1=z_1+\lambda_1} f\, d^{l-2}t.$$

Again, we move the contour of integration in the first summand to

$$\{(t_2,\dots,t_l) \in \mathbb{C}^{l-1} \,|\, t_j \in \mathcal{C}_{(\min\{-\mathrm{Re}\,(2\Lambda_1-\lambda_1^0)-\epsilon,0\},u_2^j,\dots,u_n^j)},\ j=2,\dots,l\}$$

without meeting any poles of the integrand, and the integrand of the second summand has simple poles at $t_j = z_1+\lambda_1-2$ coming from the poles at $t_2-t_j=1$. It may also have poles at $2\lambda_1=1$ coming from $t_2=z_1-\lambda_1$.

Continuing the process we prove Lemma 32. □

Lemma 33. *The analytic continuation of the function* $I(w,W)(z,\lambda,p)$ *from* z^0,λ^0,p^0 *to a neighborhood of the point* z^0,Λ,p^0 *is given by*

$$I(w,W)(z^0,\lambda,p^0) = \sum_{\bar k} \frac{l!}{n!\,k_1!\dots k_n!(l-k)!} \int\limits_{\substack{t_i\in\mathcal{C},\\ i=k+1,\dots,l}} \mathrm{res}_{\bar k} f(t,z^0,\lambda,p^0)\, d^{l-k}t, \qquad (30)$$

where $k=k_1+\dots+k_n$, $d^{l-k}t = dt_{k+1}\dots dt_l$. *Here* $\mathcal{C}$ *denotes the curve* $\mathcal{C}_u$ *with* $u_m = \min\{-\mathrm{Re}\,\Lambda_m-\varepsilon,0\}$, $m=1,\dots,n$, ϵ *is a sufficiently small positive number, and the sum is over all* $\bar k \in \mathbb{Z}^n_{\geq 0}$ *such that* $k \leq l$, *and* $k_m \leq 2\mathrm{Re}\,\Lambda_m+1$ *for* $m=1,\dots,n$. *Morever if* $B(\bar l)\bigcap B(\bar m)=\emptyset$, *then sum (30) is well defined at* z^0,Λ,p^0.

Proof: We analytically continue each summand in (29) from z^0,λ^0,p^0 to z^0,Λ,p^0. By Lemma 32, the poles of the integrand of a summand and the contour of integration can be pictured as follows.



Here $z_j = z_j^0$, $\lambda_j = \lambda_j^0$, $j = 1, \dots, n$.

Note that there are also poles of the integrand of the type $t_i = z_j^0 \pm (\lambda_j + p^0 s)$, $s = 1, 2, \dots$. These poles are far away from our picture according to our choice of p^0. Note also that all poles are simple.

We move the parameters λ_m from λ_m^0 to Λ_m. The contour of integration of each summand in Lemma 32, $\{t \in \mathbb{C}^l \,|\, t_j \in \mathcal{C} = \mathcal{C}_{u(\lambda)},\ j = k+1, \dots, l\}$, depends on λ. At every moment of deformation of parameters λ_m, we define $I(w, W)(z^0, \lambda, p^0)$ by

$$I(w,W)(z^0,\lambda,p^0) = \sum_{\bar{k}} \frac{l!}{n!\,k_1!\dots k_n!(l-k)!} \int\limits_{\substack{t_i\in\mathcal{C}_{u(\lambda)},\\ i=k+1,\dots,l}} \mathrm{res}_{\bar{k}} f(t,z^0,\lambda,p^0)\, d^{l-k}t,$$

where $u(\lambda) = (u_1(\lambda_1), \dots, u_n(\lambda_n)) \in \mathbb{R}^n$, $u_m(\lambda_m) = \min\{-\mathrm{Re}\,(2\Lambda_m - \lambda_m) - \epsilon, 0\}$, $m = 1, \dots, n$, and the sum is over all $\bar{k} \in \mathbb{Z}^n_{\geq 0}$ such that $k \leq l$, and $k_m \leq 2\mathrm{Re}\,\Lambda_m$ for $m = 1, \dots, n$.

If $\Lambda_m \notin \Lambda^+$ for all $m = 1, \dots, n$, then at every moment of the deformation the integrand is holomorphic by Lemma 31. On the drawing, the points $z_i - \lambda_i$, $i = 1, \dots, n$, move to the left and the rest of the picture moves to the right. Note that the points $z_i - \lambda_i$, $i = 1, \dots, n$, encounter neither other points on the picture nor the curve of integration $\mathcal{C}_{u(\lambda)}$.

If, for some $i \in \{1, \dots, n\}$, $\Lambda_i \in \Lambda^+$, then by Lemma 31, the function $\mathrm{res}_{\bar{k}}(z, \lambda, p)$ can have a pole at $\lambda_i = \Lambda_i$ for $\bar{k}$ such that $k_i = 2\Lambda_i$. In this case, in the picture, in the last moment of the deformation the point $z_i - \lambda_i$ coincides with the point $z_i + \lambda_i - k$.

However, if $w = w_{\bar{l}}$, $W = W_{\bar{m}}$ and $B(\bar{l}) \bigcap B(\bar{m}) = \emptyset$, then the pole is trivial by the following two lemmas.

Lemma 34. *Let* $i \in \{1, \dots, n\}$ *and* $\Lambda_i \in \Lambda^+$. *Let* $W = W_{\bar{m}}$ *and* $m_i \leq 2\Lambda_i$. *Then for any* $w = w_{\bar{l}}$,

$$\mathrm{res}_{\lambda_i=\Lambda_i}\mathrm{res}_{t_{2\Lambda_i+1}=z_i+\lambda_i-2\Lambda_i}\dots\mathrm{res}_{t_2=z_i+\lambda_i-1}\mathrm{res}_{t_1=z_i+\lambda_i}\Phi_l wW = 0.$$

Proof: Let i=1. We prove that

$$\mathrm{res}_{t_{2\Lambda_1+1}=z_1+\lambda_1-2\Lambda_1}\dots\mathrm{res}_{t_2=z_1+\lambda_1-1}\mathrm{res}_{t_1=z_1+\lambda_1}\Phi_l wW=0.$$

Recall that W has the form $W=\sum\limits_{\sigma\in\mathbb{S}^l}[\dots]_\sigma^{trig}$. The residue is nontrivial only for the terms corresponding to permutations σ, which satisfy the condition:

$$m_1\geq(\sigma^{-1})_1>(\sigma^{-1})_2>\dots>(\sigma^{-1})_{2\Lambda+1}.$$

If $m_1\leq 2\Lambda$ then there are no such permutations.

The cases $i=2,\dots,n$ are proved similarly. □

Lemma 35. *Let $i\in\{1,\dots,n\}$ and $\Lambda_i\in\Lambda^+$. Let $w=w_{\bar l}$ and $l_i\leq 2\Lambda_i$. Then for any $W=W_{\bar m}$,*

$$\mathrm{res}_{\lambda_i=\Lambda_i}\mathrm{res}_{t_{2\Lambda_i+1}=z_i+\lambda_i-2\Lambda_i}\dots\mathrm{res}_{t_2=z_i+\lambda_i-1}\mathrm{res}_{t_1=z_i+\lambda_i}\Phi_l w_{\bar l}W_{\bar m}=0.$$

Proof: Let i=1. We prove that

$$\mathrm{res}_{t_{2\Lambda_1+1}=z_1+\lambda_1-2\Lambda_1}\dots\mathrm{res}_{t_2=z_1+\lambda_1-1}\mathrm{res}_{t_1=z_1+\lambda_1}\Phi_l w_{\bar l}\tilde{W_{\bar m}}=0.$$

Recall that $w_{\bar l}$ has the form $w_{\bar l}=\sum\limits_{\sigma\in\mathbb{S}^l}[\dots]_\sigma^{rat}$. The resudue is nontrivial only for the terms corresponding to permutations σ which satisfy the condition:

$$l_1\geq(\sigma^{-1})_1>(\sigma^{-1})_2>\dots>(\sigma^{-1})_{2\Lambda+1}.$$

Since $1\notin B(\bar l)$, we have $l_1\leq 2\Lambda$ and there are no such permutations.

The cases $i=2,\dots,n$ are proved similarly. □

Lemma 33 is proved. □

Note that a residue of a meromorphic function can be described as an integral over a small circle.

For $B\in\mathbb{C}$ and $\epsilon>0$, denote $\mathcal{D}_B\subset\mathbb{C}$ the circle with center B and radius ϵ. Denote $\mathcal{D}$ the curve $\mathcal{C}_u$ with $u_m=\min\{-\mathrm{Re}\,\Lambda_m-\varepsilon,0\}$.

Lemma 36. *The analytic continuation of the function $I(w,W)$ from z^0,Λ,p^0 to z^0,Λ,P is well defined. Moreover, for sufficiently small $\epsilon>0$,*

$$I(w,W)(z^0,\Lambda,P)= \tag{31}$$
$$\sum_{m=0}^{l}\sum_{a_1,\dots,a_m}\frac{l!}{(l-m)!}\int\limits_{\substack{t_i\in\mathcal{D},\\ i=m+1,\dots,l}}\left(\sum_{\sigma\in\mathbb{S}^m}\int\limits_{\substack{t_i\in\mathcal{D}_{a_i},\\ i=1,\dots,m}}f(t,z^0,\Lambda,P)\,dt_{\sigma_1}\dots dt_{\sigma_m}\right)dt_{m+1}\dots dt_l,$$

where the second sum is over all $a=(a_1,\dots,a_m)\in\mathbb{C}^m$ such that for each $i=1,\dots,m$, there exist $j\in\{1,\dots,n\}$, $k,s\in\mathbb{Z}_{>0}$, such that $a_i=z_j+\Lambda_j-k+sP$; a_i is located to the right from $\mathcal{D}$ and $a_i\neq a_k$ for all $k,i=1,\dots,m$, $k\neq i$.

Note that formula (31) is a generalization of formula (30), however, in sum (31) there are many additional zero terms.

Proof: We analytically continue each summand in (30) with respect to p from p^0 to P preserving z^0, Λ. We move p and preserve the contour of integration $\mathcal{I}$ as long as the integrand is holomorphic for all $t \in \mathcal{I}$ at z, Λ, p (cf. Lemma 31). When a pole of the integrand goes through the contour of integration, we do the same procedure as in the proof of Lemma 32.

An individual integral on the right hand side of (31) is over the cycle

$$\{t \,|\, t_i \in \mathcal{D}_{a_i},\ i = 1, \dots, m;\ t_i \in \mathcal{D},\ i = m+1, \dots, l\}.$$

The points a_i are the poles of the integrand which were on the left of $\mathcal{D}$ at the beginning of the deformation and are on the right of $\mathcal{D}$ at the final moment of the deformation. All poles of the integrand which were on the right of $\mathcal{D}$ at the beginning of the deformation, remain on the right at the every moment of the deformation. The new poles a_i on the right of $\mathcal{D}$ do not coincide with the old poles on the right of $\mathcal{D}$ due to conditions (9)-(11) on P and Λ.

Note, that the procedure of analytic continuation, described in Lemma 32, gives decomposition (31) in which the sum is over $a = (a_1, \dots, a_m)$, where some of coordinates a_i could be equal. However, such integrals are equal to zero. In fact, let $a_i = a_j$ for some $i < j$. When we integrate over the variables $t_1, \dots, t_{j-1}$, the resulting integrand could have a pole at $t_j = a_j$ of at most first order. Then the reason of Lemma 30 shows that the resulting integrand is holomorphic at $t_j = a_j$. $\square$

Let, $\mathcal{D}'$ be the line $\{x \in \mathbb{C} \,|\, \operatorname{Re} x = \min\{\operatorname{Re}(Z_i - \Lambda_i),\ i = 1, \dots, n\} - \epsilon\}$.

Lemma 37. *The analytic continuation from z^0, Λ, P to Z, Λ, P is well defined. Moreover, for sufficiently small $\epsilon > 0$,*

$$I(w, W)(Z, \Lambda, P) = \tag{32}$$

$$\sum_{m=0}^{l} \sum_{b_1, \dots, b_m} \frac{l!}{(l-m)!} \int\limits_{\substack{t_i \in \mathcal{D}', \\ i = m+1, \dots, l}} \left(\sum_{\sigma \in \mathbb{S}^m} \int\limits_{\substack{t_i \in \mathcal{D}_{b_i}, \\ i = 1, \dots, m}} f(t, Z, \Lambda, P)\, dt_{\sigma_1} \dots dt_{\sigma_m} \right) dt_{m+1} \dots, dt_l,$$

where the second sum is over all $b = (b_1, \dots, b_m) \in \mathbb{C}^m$ such that for each $i = 1, \dots, m$, there exist $j \in \{1, \dots, n\}$, $k, s \in \mathbb{Z}_{>0}$, such that $b_i = Z_j + \Lambda_j - k + sP$; and b_i is located to the right from $\mathcal{D}'$.

Proof:

First, we move the parameters z_m from z_m^0 to $z_m^0 + \operatorname{Re} Z_m$. The contour of integration of a summand in (31) has the form

$$\{t \in \mathbb{C}^l \,|\, t_i \in \mathcal{D}_{a_i(z)},\ i = 1, \dots, m;\ t_i \in \mathcal{D} = \mathcal{C}_{u(z)},\ i = m+1, \dots, l\},$$

where $u(z) = \mathrm{Re}\,(z_m - \Lambda_m) - \epsilon$, and depends on z. At every moment of the deformation of parameters z_m, we define $I(w,W)(z,\Lambda,P)$ by

$$I(w,W)(z,\Lambda,P) =$$

$$\sum_{m=0}^{l} \sum_{a_1,\dots,a_m} \frac{l!}{(l-m)!} \int\limits_{\substack{t_i \in \mathcal{C}_{u(z)},\\ i=m+1,\dots,l}} \left(\sum_{\sigma \in \mathbb{S}^m} \int\limits_{\substack{t_i \in \mathcal{D}_{a_i},\\ i=1,\dots,m}} f(t,z,\Lambda,P)\, dt_{\sigma_1} \dots dt_{\sigma_m} \right) dt_{m+1} \dots , dt_l,$$

where the second sum is over all $a = (a_1,\dots,a_m) \in \mathbb{C}^m$ such that for each $i = 1,\dots,m$, there exist $j \in \{1,\dots,n\}$, $k, s \in \mathbb{Z}_{>0}$, such that $a_i = z_j + \Lambda_j - k + sP$; and a_i is located to the right from $\mathcal{C}_{u(z)}$.

Consider the summand related to $a \in \mathbb{C}^m$. We move the contour of integration with respect to $t_{m+1},\dots,t_l$ from $\mathcal{C}_{u(z^0 + \mathrm{Re}\, Z)}$ to $\mathcal{D}'$. We use the method described in Lemma 32.

Finally, we move parameters $z_1 \dots, z_n$ from $z_1^0 + \mathrm{Re}\, Z_1, \dots, z_n^0 + \mathrm{Re}\, Z_n$ to $Z_1 \dots, Z_n$ in the same way as we moved $\lambda_1,\dots,\lambda_n$ in Lemma 33.

An individual integral in the RHS of (32) is over the cycle

$$\{t \,|\, t_i \in \mathcal{D}_{b_i},\ i = 1,\dots,m;\ t_i \in \mathcal{D}',\ i = m+1,\dots,l\}.$$

The points b_i are the poles of the integrand which were on the left of $\mathcal{D}$ at the beginning of the deformation and are on the right of $\mathcal{D}'$ at the final moment of the deformation. All poles of the integrand which were on the right of $\mathcal{D}$ at the beginning of the deformation, remain on the right of $\mathcal{D}'$ at the every moment of the deformation. The new poles b_i on the right of $\mathcal{D}'$ do not coincide with the old poles on the right of $\mathcal{D}'$ due to conditions (12) on Z. □

Theorem 10 is proved. □

5.2. **Proof of Theorems 11, 12.** Let $\mathrm{Im}\,\mu \neq 0$. Fix a rational weight function $w = w_{\bar l}$, and a trigonometric weight function $W = W_{\bar m} \in \mathfrak{S}$. Fix $\Lambda \in \mathbb{C}^n$.

We fix parameters z, p, λ^0 in such a way that $\mathrm{Re}\,\lambda_m^0 < 0$, p is a negative number with large absolute value and $|\mathrm{Im}\,(z_k + \Lambda_k - z_m - \Lambda_m)|$ is large for $k, m = 1,\dots,n$, $k \neq m$. Namely, let $p = -2 \sum\limits_{j=1}^{n} |\Lambda_j| - 1$, $\lambda_k^0 = -1 + i\mathrm{Im}\,\Lambda_k$, $z_k^0 = i\,(-\mathrm{Im}\,\Lambda_k + 3kA)$, where A is a large real number such that $A > 2|\Lambda_k|$, $k = 1,\dots,n$.

By Lemma 33,

$$I(w,W)(z,\lambda,p) = \sum_{\bar k} \frac{l!}{n!\, k_1! \dots k_n! (l-k)!} \int\limits_{\substack{t_i \in \mathcal{C},\\ i=k+1,\dots,l}} \mathrm{res}_{\bar k} f(t,z,\lambda,p)\, d^{l-k}t, \qquad (33)$$

where $k = k_1 + \dots + k_n$, $\mathcal{C}$ denotes the curve $\mathcal{C}_u$ with $u_m = \min\{-\mathrm{Re}\,(2\Lambda_m - \lambda_m) - \varepsilon, 0\}$, $m = 1,\dots,n$, ϵ is a sufficiently small positive number, and the sum is over all $\bar k \in \mathbb{Z}^n_{\geq 0}$ such that $k \leq l$, and $k_m \leq 2\mathrm{Re}\,\Lambda_m + 1$ for $m = 1,\dots,n$.

Let $i \in \{1,\dots,n\}$. We have the following cases.

1). If i does not belong to $B_\Lambda(\bar m)$, then the analytic continuation of the function $I(w,W)(z,\lambda,p)$ with respect to λ_i from λ_i^0 to Λ_i is well defined, see the proof of Lemma 33.

2). If i belongs to $B_\Lambda(\bar m)$ and i does not belong to $B_\Lambda(\bar l)$, then the analytic continuation of the function $I(w,W)(z,\lambda,p)$ with respect to λ_i from λ_i^0 to Λ_i is well defined, see the proof of Lemma 33. In this case $c_{\bar m}(\lambda,p)$ is equal to zero at $\lambda_i=\Lambda_i$, so $J_{\bar l,\bar m}(z,\lambda,p)=c_{\bar m}(\lambda,p)I(w,W)(z,\lambda,p)$ is equal to zero at $\lambda_i=\Lambda_i$.

3). If i belongs to $B_\Lambda(\bar m)$ and to $B_\Lambda(\bar l)$, then in decomposition (33) all the summands are well defined at $\lambda_i=\Lambda_i$ except for the summands corresponding to $\bar k$ such that $k_i=2\Lambda_i+1$. Such summands have a simple pole at $\lambda_i=\Lambda_i$, see the proof of Lemma 33. In this case $c_{\bar m}(\lambda,p)$ is equal to zero at $\lambda_i=\Lambda_i$, so the function $J_{\bar l,\bar m}(z,\lambda,p)=c_{\bar m}(\lambda,p)I(w,W)(z,\lambda,p)$ is well defined at $\lambda_i=\Lambda_i$.

This proves Theorem 11.

Moreover, if $B_\Lambda(\bar l)=B_\Lambda(\bar m)=B$, then we have

$$J_{\bar l,\bar m}(z,\Lambda,p)=\sum_{\bar k}\frac{l!}{n!\,k_1!\dots k_n!(l-k)!}\int\limits_{\substack{t_i\in\mathcal{C},\\ i=k+1,\dots,l}}(c_{\bar m}\mathrm{res}_{\bar k}f)(t_{k+1},\dots,t_l,z,\Lambda,p)\,d^{l-k}t,$$

where $\mathcal{C}$ denotes the curve $\mathcal{C}_u$ with $u_m=\min\{-\mathrm{Re}\,\Lambda_m-\varepsilon,0\}$, $m=1,\dots,n$, and the sum is over all $\bar k\in\mathbb{Z}^n_{\geq 0}$ such that $k\leq l$, $k_m\leq 2\mathrm{Re}\,\Lambda_m+1$ for $m=1,\dots,n$, and $k_m=2\Lambda_m+1$ for all $m\in B$.

Let $j\in B$. We compute the residues at points $z_j+\lambda_j,z_j+\lambda_j-1,\dots,z_j+\lambda_j-2\Lambda_j$ explicitly using the following Lemma.

Lemma 38. *Let $B_\Lambda(\bar l)=B_\Lambda(\bar m)=B$ and let $j\in B$, so that $\Lambda_j\in\Lambda^+$, $\bar l,\bar m\in\mathcal{Z}^n_l$ and $l_j,m_j>2\Lambda_j$. Let $k=2\Lambda_j$,*

$$\begin{aligned}&(c_{\bar m}\mathrm{res}_{t_{k+1}=z_j+\lambda_j-k}\dots\mathrm{res}_{t_2=z_j+\lambda_j-1}\mathrm{res}_{t_1=z_j+\lambda_j}\Phi_l w_{\bar l}W_{\bar m})(t_{k+2},\dots,t_l,z,\Lambda)=\\&\qquad(k+1)!\,e^{\mu(k+1)z_j\pi i/p}\,\psi_{m_j,k}\,\varphi_{\Lambda,j,k}(z)\,(c_{\bar m^{(j)}}\Phi_{l-k-1}w_{\bar l^{(j)}}W_{\bar m^{(j)}})(t_{k+2},\dots,t_l,z,\Lambda^{(j)}),\end{aligned}$$

where $\bar l^{(j)}=(l_1,\dots,l_{j-1},l_j-k-1,l_{j+1},\dots,l_n)$, $\bar m^{(j)}=(m_1,\dots,m_{j-1},m_j-k-1,m_{j+1},\dots,m_n)$ and $\bar\Lambda^{(j)}=(\Lambda_1,\dots,\Lambda_{j-1},-\Lambda_j-1,\Lambda_{j+1},\dots,\Lambda_n)$.

Proof: Consider the case $j=1$. For a function $g(t,z,\lambda)$, the coefficient in the Laurent expansion with respect to variables $t_1,...,t_{2\Lambda_1+1}$ at $t_i=z_1+\lambda_1-i+1$, $i=1,\dots,2\Lambda_1+1$, computed at $\lambda_1=\Lambda_1$, i.e. the coefficient of

$$\prod_{i=1}^{2\Lambda_1+1}\frac{1}{t_i-z_1-\lambda_1+i-1},$$

is called the *main coefficient*. We compute the main coefficients of the functions $\Phi_l,w_{\bar l},W_{\bar m}$.

The main coefficient of the function $\Phi_l(t,z,\lambda)$ is

$$\begin{aligned}&e^{\mu(2\Lambda_1+1)z_j\pi i/p}\,a_1(\Lambda)\,\varphi_{\Lambda^{(1)},1,2\Lambda_1}(z)\prod_{j=2\Lambda_1+2}^{l}\prod_{i=1}^{2\Lambda_1+1}\frac{\Gamma((z_1+\Lambda_1-i+1-t_j-1)/p)}{\Gamma((z_1+\Lambda_1-i+1-t_j+1)/p)}\times\\&\times\prod_{j=2\Lambda_1+2}^{l}\prod_{i=1}^{n}\frac{\Gamma((t_j-z_i+\lambda_i)/p)}{\Gamma((t_j-z_i-\lambda_i)/p)}\prod_{2\Lambda_1+1<i<j\leq l}\frac{\Gamma((t_i-t_j-1)/p)}{\Gamma((t_i-t_j+1)/p)}.\end{aligned}$$

Here $a_1(\Lambda_1) \in \mathbb{C}$ is some explicitly computable constant. Note that $\varphi_{\Lambda^{(1)},1,2\Lambda_1}(z) = \varphi_{\Lambda,1,2\Lambda_1}(z)$. The main coefficient of the function $\Phi_l(t,z,\lambda)$ is equal to

$$\begin{gathered}\prod_{j=2\Lambda_1+2}^{l} \frac{t_j - z_1 - \Lambda_1}{t_j - z_1 + \Lambda_1}\, \frac{t_j - z_1 + \Lambda'_1}{t_j - z_1 - \Lambda'_1}\, \frac{\sin((t_j - z_1 - \Lambda_1)\pi i/p)}{\sin((t_j - z_1 + \Lambda_1)\pi i/p)}\, \frac{\sin((t_j - z_1 + \Lambda'_1)\pi i/p)}{\sin((t_j - z_1 - \Lambda'_1)\pi i/p)} \times \\ \times a_1(\Lambda)\, e^{\mu(2\Lambda_1+1)z_j\pi i/p}\, \varphi_{\Lambda,1,2\Lambda_1}(z)\, \Phi_{l-2\Lambda_1-1}(t_{2\Lambda_1+1}, \dots, t_l, z, \Lambda^{(1)}),\end{gathered}$$

where $\Lambda'_1 = -\Lambda_1 - 1$.

The function $w_{\bar{l}}(t,z,\lambda)$ has the form $w_{\bar{l}} = \sum_{\sigma \in \mathbb{S}^l} [\dots]^{rat}_{\sigma}$. Consider the sum over $\sigma \in \mathbb{S}^l$ such that $\sigma(i) = i$ for all $i = l_1 + 1, ..., l$. We have

$$\begin{aligned} &w_{\bar{l}}(t,z,\lambda) = a_2(\Lambda_1) \times \\ &\times \sum_{\sigma \in \mathbb{S}^l} \left[\prod_{1 \le i < j \le l_1} \frac{t_i - t_j}{t_i - t_j + 1} \prod_{i=1}^{l} \frac{1}{t_i - z_1 - \lambda_1} \prod_{i=l_1+1}^{l} (t_j - z_1 + \lambda_1) w_{(l_2,\dots,l_n)}(t_{l_1+1}, \dots, t_l, z, \lambda) \right]^{rat}_{\sigma}. \end{aligned}$$

Here, again, $a_2(\Lambda_1) \in \mathbb{C}$ is an easily computable constant.

Let $\sigma \in \mathbb{S}^l$ be a permutation and there exists $i \le 2\Lambda_1 + 1$ such that $\sigma(i) > l_1$. Then the term corresponding to σ does not contribute to the main coefficient. Hence, this coefficient is

$$\begin{aligned} \tilde{a}_2(\Lambda_1) \sum_{\sigma \in \mathbb{S}^{l-2\Lambda_1-1}} \Bigg[& \prod_{j=2\Lambda_1+2}^{l_1} \prod_{i=1}^{2\Lambda+1} \frac{z_1 + \Lambda_1 - i + 1 - t_j - 1}{z_1 + \Lambda_1 - i + 1 - t_j + 1} \prod_{i=2\Lambda_1+2}^{l} \frac{1}{t_i - z_1 - \lambda_1} \times \\ \times & \prod_{i=l_1+1}^{l} (t_j - z_1 + \lambda_1) \prod_{2\Lambda_1+2 \le i < j \le l_1} \frac{t_i - t_j}{t_i - t_j + 1} w_{(l_2,\dots,l_n)}(t_{l_1+1}, \dots, t_l, z, \Lambda_1, \lambda_2, \dots, \lambda_n) \Bigg]^{rat}_{\sigma}, \end{aligned}$$

where $\tilde{a}_2(\Lambda_1) \in \mathbb{C}$ is another easily computable constant and the group $\mathbb{S}^{l-2\Lambda_1-1}$ permutes the variables $t_{2\Lambda_1+1}, \dots, t_l$. Simplifying, we get the main coefficient of the function $w_{\bar{l}}(t,z,\lambda)$ is given by

$$\tilde{a}_2(\Lambda_1) \prod_{j=2\Lambda+1}^{l} \frac{t_j - z_1 + \Lambda_1}{t_j - z_1 - \Lambda_1}\, \frac{t_j - z_1 - \Lambda'_1}{t_j - z_1 + \Lambda'_1}\, w_{(l'_1, l_2 \dots, l_n)}.$$

The main coefficient of the function $W_{\bar{m}}(t,z,\lambda)$ is computed similarly and is equal to

$$a_3(\Lambda_1) \prod_{j=2\Lambda_1+1}^{l} \frac{\sin((t_j - z_1 + \Lambda_1)\pi i/p)}{\sin((t_j - z_1 - \Lambda_1)\pi i/p)}\, \frac{\sin((t_j - z_1 - \Lambda'_1)\pi i/p)}{\sin((t_j - z_1 + \Lambda'_1)\pi i/p)}\, W_{(m'_1, m_2 \dots, m_n)},$$

where $a_3(\Lambda_1) \in \mathbb{C}$ is some easily computable comstant.

Multiplying the above main coefficients we get the statement of the Lemma. We have

$$a_1(\Lambda_1)\tilde{a}_2(\Lambda_1)a_3(\Lambda_1)c_{\bar{m}}(\Lambda) = (k+1)!\, c^{(1)}_{\bar{m}}(\Lambda^{(1)})\, \psi_{2\Lambda_1}.$$

The cases $j = 2, \dots, n$ are proved similarly. □

We have

$$J_{\bar l,\bar m}(z,\Lambda,p)=\sum_{\bar k}\frac{l!}{n!\,k_1!\dots k_n!(l-k)!}\times \tag{34}$$

$$\times\int\limits_{\substack{t_i\in\mathcal{C},\\ i=k+1+l'(B),\dots,l}}\mathfrak{C}_{\Lambda,\bar m}(c_{\bar m'(B)}\mathrm{res}_{\bar k}\Phi_{l'(B)}w_{\bar l'(B)}W_{\bar m'(B)})(t_{k+1+l'(B)},\dots,t_l,z,\Lambda,p)\,d^{l-k-l'(B)}t,$$

where $\mathcal{C}$ denotes the curve $\mathcal{C}_u$ with $u_m=\min\{-\mathrm{Re}\,\Lambda_m-\varepsilon,0\}$, $m=1,\dots,n$, and the sum is over all $\bar k\in\mathbb{Z}^n_{\geq 0}$ such that $k\leq l-l'(B)$, $k_m\leq 2\mathrm{Re}\,\Lambda_m+1$ for $m=1,\dots,n$, and $k_m=0$ for all $m\in B$.

Let $\mathcal{C}'$ be the curve $\mathcal{C}_u$ with $u_m=\min\{-\mathrm{Re}\,\Lambda'_m-\varepsilon,0\}$, $m=1,\dots,n$. Then the integrand of each integral in sum (34) does not have poles of the type $t_j=a$, where $a\in\mathbb{C}$ is located between $\mathcal{C}'$ and $\mathcal{C}$, see Lemma 31. Thus, we can move the contour of integration in each summand to $\{t\in\mathbb{C}^{l-k-l'(B)}\,|\,t_i\in\mathcal{C}',\ i=k+1+l'(B),\dots,l\}$.

We proved Theorem 12 for our choice of z,p. Theorem 12 holds for all z,p, since both left and right hand sides of the formula of Theorem 12 are meromorphic functions of parameters z,p. □

5.3. **Proof of Theorem 19.** Consider a matrix

$$I^l(z,\lambda,p)=\{\prod_{i\in B_\Lambda(\bar m)}(\lambda_i-\Lambda_i)I_{\bar l,\bar m}(z,\lambda,p)\}_{\bar l,\bar m\in\mathcal{Z}^n_l}.$$

The matrix $I^l(z,\lambda,p)$ is obtained from the matrix $J^l(z,\lambda)=J^l(z,\lambda,p)$ by multiplication the $\bar m$-th row by $(\prod_{i\in B_\Lambda(\bar m)}(\lambda_i-\Lambda_i))/c_{\bar m}(\lambda)$ for all $\bar m\in\mathcal{Z}^n_l$. For generic p, these factors are well defined and not equal to zero. All entries of the matrix $I^l(z,\lambda,p)$ are well defined and the matrix $I^l(z,\lambda)$ is upper block triangular at $\lambda=\Lambda$.

It follows from formula 5.14 in [TV1] that the matrix $I^l(z,\lambda)$ is non-degenerate for all z,λ,p satisfying conditions (9)-(14).

Hence, the matrix

$$I^l_{\mathrm{adm}}(z,\lambda,p)=\{\prod_{i\in B_\Lambda(\bar m)}(\lambda_i-\Lambda_i)I_{\bar l,\bar m}(z,\lambda,p)\},$$

where $\bar l,\bar m\in\mathcal{Z}^n_l$ run through the set of all Λ-admissible indices, is also non-degenerate for all z,λ,p satisfying conditions (9)-(14).

For all p,Λ satisfying conditions (9)-(14) and Λ-admissible indices $\bar m$, the factors $(\prod_{i\in B_\Lambda(\bar m)}(\lambda_i-\Lambda_i))/c_{\bar m}(\lambda)$ are well defined and non-zero. Hence, the matrix $J^l_{\mathrm{adm}}(z,\lambda,p)$ is also non-degenerate for all z,λ,p satisfying conditions (9)-(14).

The matrix $J^l(z,\lambda)$ is upper block triangular at $\lambda=\Lambda$. Now, formula (21) follows from the Inclusion-Exclusion Principle and Theorem 12.

The Theorem is proved. □

5.4. **Proof of Theorem 1.** The rational R-matrix $R_{\lambda_1\lambda_2}(x)\in\mathrm{End}(V_{\lambda_1}\otimes V_{\lambda_2})$ is a meromorphic function of λ_1,λ_2, defined explicitly in Section 2.3 for $\lambda_1,\lambda_2\notin\Lambda^+$.

Consider the qKZ equation with $n = 2$, $\mu = 2$, $p = -2(|\lambda_1| + |\lambda_2|) - 1$. Consider the functions $\{\Psi_{\bar m}(z, \lambda) \,|\, \bar m \in \mathcal{Z}_l^2\}$ given by

$$\Psi_{\bar m}(z, \lambda) = \sum_{l_1 + l_2 = l} J_{\bar l, \bar m}(z, \lambda)\, f^{l_1} v_1 \otimes f^{l_2} v_2,$$

where the function $J_{\bar l, \bar m}$ is defined in (15).

The functions $\{\Psi_{\bar m}(z, \lambda) \,|\, \bar m \in \mathcal{Z}_l^2\}$ are solutions of the qKZ equation. Moreover, for generic λ, the values of these functions span $V_{\lambda_1} \otimes V_{\lambda_2}$ over $\mathbb{C}$ by Theorem 5.14 in [TV1]. By Theorem 11, these functions are well defined for all λ_1, λ_2. Moreover, they span $V_{\lambda_1} \otimes V_{\lambda_2}$, for all λ_1, λ_2 and generic z, see theorem 5.14 in [TV1] and Section 4.4. The first statement of Theorem 1 follows, now, from the equality of meromorphic functions:

$$\Psi_{\bar m}(z_1 + p, z_2, \lambda_1, \lambda_2) = e^{-\mu h_1} R_{\lambda_1 \lambda_2}(z_1 - z_2) \Psi_{\bar m}(z_1, z_2, \lambda_1, \lambda_2)$$

for all $\bar m \in \mathcal{Z}_l^2$.

For all complex λ_1, λ_2, the R-matrix $R_{\lambda_1 \lambda_2}(x)$ commutes with the action of sl_2.

If $\lambda_1, \lambda_2 \notin \Lambda^+$, then $S_{\lambda_1} \otimes V_{\lambda_2} + V_{\lambda_1} \otimes S_{\lambda_2}$ is the zero submodule and the second statement of the Theorem is trivial. Otherwise, let $\lambda \in \{\lambda_1, \lambda_2\}$ be the minimal dominant weight. Namely, if $\lambda_1 \in \Lambda^+$ and $\lambda_2 \notin \Lambda^+$, set $\lambda = \lambda_1$. If $\lambda_2 \in \Lambda^+$ and $\lambda_1 \notin \Lambda^+$, set $\lambda = \lambda_2$. If $\lambda_1 \in \Lambda^+$ and $\lambda_2 \in \Lambda^+$, set $\lambda = \min\{\lambda_1, \lambda_2\}$. Then the submodule $S_{\lambda_1} \otimes V_{\lambda_2} + V_{\lambda_1} \otimes S_{\lambda_2}$ is generated by all singular vectors in $V_{\lambda_1} \otimes V_{\lambda_2}$ of weights less than $\lambda_1 + \lambda_2 - 2\lambda$. Hence, the R-matrix $R_{\lambda_1 \lambda_2}(x)$ preserves $S_{\lambda_1} \otimes V_{\lambda_2} + V_{\lambda_1} \otimes S_{\lambda_2}$.

The third statement of the Theorem follows from the first two statements. □

References

[CP] V.Chari and A.Pressley, *A guide to quantum groups*,Cambrige University Press, Cambrige, 1994.

[F1] G.Felder, *Conformal field theory and integrable systems associated to elliptic curves*, in Proceedings of the ICMP, Zurich, 1994, Birkhauser, 1994.

[F2] G.Felder, *Elliptic quantum groups*, in Proceedings of the ICMP, Paris, 1994 (D.Iagolnitzer, ed.), Intern. Press, Boston, 1995, 211-218.

[FR] I. Frenkel and N. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Commun. Math. Phys. 146 (1992), 1–60

[FTV1] G.Felder, V.Tarasov and A.Varchenko, *Solutions of elliptic qKZB equations and Bethe ansatz I*, in Topics in Singularity Theory, V.I.Arnold's 60th Aniversary Collection, Advances in the Mathematical Sciences -34, AMS Translations, Series 2, v.180, 45-76.

[FTV2] G.Felder, V.Tarasov and A.Varchenko, *Monodromy of solutions of the elliptic quantum Knizhnik-Zamolodchikov-Bernard difference equation*, Preprint (1997).

[FV] G.Felder and A.Varchenko,*On representations of the elliptic quantum group $E_{\tau,\eta}(sl_2)$*, Comm.Math.Phys. 181, (1996), 741-461.

[JM] M.Jimbo and T.Miwa, *Algebraic analysis of solvable lattice models*, CBMS Regional Conf.Series in Math, 85 (1995).

[KRS] P.P.Kulish, N.Yu.Reshetikhin and E.K.Sklyanin, *Yang-Baxter equation and representation theory I*, Lett.Math.Phys. 5 (1981), 393-403.

[S] F.A. Smirnov, *Form factors in completely integrable models of quantum field theory*, Advanced Series in Math. Phys., vol. 14, World Scientific, Singapore, 1992.

[T] V.O. Tarasov, *Irreducible monodromy matrices for the R-matrix of the XXZ-model and lattice local Hamiltonians*, Theor. Math. Phys. 63 (1985), 440-454.

[TV1] V. Tarasov and A. Varchenko, *Geometry of q-hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. math. 128, (1997), 501-588.

[TV2] V. Tarasov and A. Varchenko, *Geometry of q-hypergeometric functions, quantum affine algebras and elliptic quantum groups*, preprint q-alg/9703044.

[TV3] V. Tarasov and A. Varchenko,*Asymptotic solutions to the quantized Knizhnik Zamolodchikov equation and Bethe vectors*, Amer.Math.Soc.Transl. (2),vol. 174, (1996), 235-273.